

Title: HYBRID MAIZE 33N56	Anticipated Class:	800
	Subclass:	200
Inventor(s): Robert Lee Segebart	Art Unit:	1638

FIELD OF THE INVENTION

This invention relates generally to the field of maize breeding, specifically relating to hybrid maize designated 33N56.

BACKGROUND OF THE INVENTION

Plant Breeding

The goal of plant breeding is to combine in a single variety or hybrid various desirable traits. For field crops, these traits may include resistance to diseases and insects, tolerance to heat and drought, reducing the time to crop maturity, greater yield, and better agronomic quality. With mechanical harvesting of many crops, uniformity of plant characteristics such as germination and stand establishment, growth rate, maturity, and plant and ear height is important.

Field crops are bred through techniques that take advantage of the plant's method of pollination. A plant is self-pollinated if pollen from one flower is transferred to the same or another flower of the same plant. A plant is sib pollinated when individuals within the same family or line are used for pollination. A plant is cross-pollinated if the pollen comes from a flower on a different plant from a different family or line. The term "cross pollination" and "out-cross" as used herein do not include self pollination or sib pollination.

Plants that have been self-pollinated and selected for type for many generations become homozygous at almost all gene loci and produce a uniform population of true breeding progeny. A cross between two different homozygous lines produces a uniform population of hybrid plants that may be heterozygous for

many gene loci. A cross of two plants each heterozygous at a number of gene loci will produce a population of heterogeneous plants that differ genetically and will not be uniform.

Maize (*Zea mays* L.), often referred to as corn in the United States, can be bred by both self-pollination and cross-pollination techniques. Maize has separate male and female flowers on the same plant, located on the tassel and the ear, respectively. Natural pollination occurs in maize when wind blows pollen from the tassels to the silks that protrude from the tops of the ears.

The development of a hybrid maize variety in a maize plant breeding program involves three steps: (1) the selection of plants from various germplasm pools for initial breeding crosses; (2) the selfing of the selected plants from the breeding crosses for several generations to produce a series of inbred lines, which, individually breed true and are highly uniform; and (3) crossing a selected inbred line with an unrelated inbred line to produce the hybrid progeny (F₁). After a sufficient amount of inbreeding successive filial generations will merely serve to increase seed of the developed inbred. Preferably, an inbred line should comprise homozygous alleles at about 95% or more of its loci.

During the inbreeding process in maize, the vigor of the lines decreases. Vigor is restored when two different inbred lines are crossed to produce the hybrid progeny (F₁). An important consequence of the homozygosity and homogeneity of the inbred lines is that the hybrid created by crossing a defined pair of inbreds will always be the same. Once the inbreds that create a superior hybrid have been identified, a continual supply of the hybrid seed can be produced using these inbred parents and the hybrid corn plants can then be generated from this hybrid seed supply.

Large scale commercial maize hybrid production, as it is practiced today, requires the use of some form of male sterility system which controls or inactivates male fertility. A reliable method of controlling male fertility in plants also offers the opportunity for improved plant breeding. This is especially true for development of maize hybrids, which relies upon some sort of male sterility system. There are several ways in which a maize plant can be manipulated so that is male sterile.

These include use of manual or mechanical emasculation (or detasseling), cytoplasmic genetic male sterility, nuclear genetic male sterility, gametocides and the like.

Hybrid maize seed is often produced by a male sterility system incorporating manual or mechanical detasseling. Alternate strips of two inbred varieties of maize are planted in a field, and the pollen-bearing tassels are removed from one of the inbreds (female) prior to pollen shed. Providing that there is sufficient isolation from sources of foreign maize pollen, the ears of the detasseled inbred will be fertilized only from the other inbred (male), and the resulting seed is therefore hybrid and will form hybrid plants.

The laborious detasseling process can be avoided by using cytoplasmic male-sterile (CMS) inbreds. Plants of a CMS inbred are male sterile as a result of factors resulting from the cytoplasmic, as opposed to the nuclear, genome. Thus, this characteristic is inherited exclusively through the female parent in maize plants, since only the female provides cytoplasm to the fertilized seed. CMS plants are fertilized with pollen from another inbred that is not male-sterile. Pollen from the second inbred may or may not contribute genes that make the hybrid plants male-fertile. The same hybrid seed, a portion produced from detasseled fertile maize and a portion produced using the CMS system can be blended to insure that adequate pollen loads are available for fertilization when the hybrid plants are grown.

There are several methods of conferring genetic male sterility available, such as multiple mutant genes at separate locations within the genome that confer male sterility, as disclosed in U.S. Patent Nos. 4,654,465 and 4,727,219 to Brar *et al.* and chromosomal translocations as described by Patterson in U.S. Patent Nos. 3,861,709 and 3,710,511. These and all patents referred to are incorporated by reference. In addition to these methods, Albertsen *et al.*, of Pioneer Hi-Bred, U.S. Patent No. 5,432,068, have developed a system of nuclear male sterility which includes: identifying a gene which is critical to male fertility; silencing this native gene which is critical to male fertility; removing the native promoter from the essential male fertility gene and replacing it with an inducible promoter; inserting this genetically engineered

gene back into the plant; and thus creating a plant that is male sterile because the inducible promoter is not "on" resulting in the male fertility gene not being transcribed. Fertility is restored by inducing, or turning "on", the promoter, which in turn allows the gene that confers male fertility to be transcribed.

There are many other methods of conferring genetic male sterility in the art, each with its own benefits and drawbacks. These methods use a variety of approaches such as delivering into the plant a gene encoding a cytotoxic substance associated with a male tissue specific promoter or an antisense system in which a gene critical to fertility is identified and an antisense to that gene is inserted in the plant (see: Fabinjanski, *et al.* EPO 89/3010153.8 Publication No. 329,308 and PCT Application PCT/CA90/00037 published as WO 90/08828).

Another system useful in controlling male sterility makes use of gametocides. Gametocides are not a genetic system, but rather a topical application of chemicals. These chemicals affect cells that are critical to male fertility. The application of these chemicals affects fertility in the plants only for the growing season in which the gametocide is applied (see Carlson, Glenn R., U.S. Patent No. 4,936,904). Application of the gametocide, timing of the application and genotype specificity often limit the usefulness of the approach and it is not appropriate in all situations.

The use of male sterile inbreds is but one factor in the production of maize hybrids. The development of maize hybrids in a maize plant breeding program requires, in general, the development of homozygous inbred lines, the crossing of these lines, and the evaluation of the crosses. Maize plant breeding programs combine the genetic backgrounds from two or more inbred lines or various other germplasm sources into breeding populations from which new inbred lines are developed by selfing and selection of desired phenotypes. Hybrids also can be used as a source of plant breeding material or as source populations from which to develop or derive new maize lines. Plant breeding techniques known in the art and used in a maize plant breeding program include, but are not limited to, recurrent selection, backcrossing, double haploids, pedigree breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, and

transformation. Often a combination of these techniques are used. The inbred lines derived from hybrids can be developed using plant breeding techniques as described above. New inbreds are crossed with other inbred lines and the hybrids from these crosses are evaluated to determine which of those have commercial potential.

Backcrossing can be used to improve inbred lines and a hybrid which is made using those inbreds. Backcrossing can be used to transfer a specific desirable trait from one line, the donor parent, to an inbred called the recurrent parent which has overall good agronomic characteristics yet that lacks the desirable trait. This transfer of the desirable trait into an inbred with overall good agronomic characteristics can be accomplished by first crossing a recurrent parent to a donor parent (non-recurrent parent). The progeny of this cross is then mated back to the recurrent parent followed by selection in the resultant progeny for the desired trait to be transferred from the non-recurrent parent. Typically after four or more backcross generations with selection for the desired trait, the progeny will contain essentially all genes of the recurrent parent except for the genes controlling the desired trait. But the number of backcross generations can be less if molecular markers are used during the selection or elite germplasm is used as the donor parent. The last backcross generation is then selfed to give pure breeding progeny for the gene(s) being transferred.

Backcrossing can also be used in conjunction with pedigree breeding to develop new inbred lines. For example, an F1 can be created that is backcrossed to one of its parent lines to create a BC1. Progeny are selfed and selected so that the newly developed inbred has many of the attributes of the recurrent parent and some of the desired attributes of the non-recurrent parent.

Recurrent selection is a method used in a plant breeding program to improve a population of plants. The method entails individual plants cross pollinating with each other to form progeny which are then grown. The superior progeny are then selected by any number of methods, which include individual plant, half sib progeny, full sib progeny, selfed progeny and topcrossing. The selected progeny are cross pollinated with each other to form progeny for another population. This population is planted and again superior plants are selected to cross pollinate with each other. Recurrent

selection is a cyclical process and therefore can be repeated as many times as desired. The objective of recurrent selection is to improve the traits of a population. The improved population can then be used as a source of breeding material to obtain inbred lines to be used in hybrids or used as parents for a synthetic cultivar. A synthetic cultivar is the resultant progeny formed by the intercrossing of several selected inbreds. Mass selection is a useful technique when used in conjunction with molecular marker enhanced selection.

Molecular markers including techniques such as Isozyme Electrophoresis, Restriction Fragment Length Polymorphisms (RFLPs), Randomly Amplified Polymorphic DNAs (RAPDs), Arbitrarily Primed Polymerase Chain Reaction (AP-PCR), DNA Amplification Fingerprinting (DAF), Sequence Characterized Amplified Regions (SCARs), Amplified Fragment Length Polymorphisms (AFLPs), Single Nucleotide Polymorphisms (SNPs) and Simple Sequence Repeats (SSRs) may be used in plant breeding methods utilizing 33N56. One use of molecular markers is Quantitative Trait Loci (QTL) mapping. QTL mapping is the use of markers, which are closely linked to alleles that have measurable effects on a quantitative trait. Selection in the breeding process is based upon the accumulation of markers linked to the positive effecting alleles and/or the elimination of the markers linked to the negative effecting alleles from the plant's genome.

Molecular markers can also be used during the breeding process for the selection of qualitative traits. For example, markers closely linked to alleles or markers containing sequences within the actual alleles of interest can be used to select plants that contain the alleles of interest during a backcrossing breeding program. The markers can also be used to select for the genome of the recurrent parent and against the markers of the donor parent. Using this procedure can minimize the amount of genome from the donor parent that remains in the selected plants. It can also be used to reduce the number of crosses back to the recurrent parent needed in a backcrossing program. The use of molecular markers in the selection process is often called Genetic Marker Enhanced Selection.

The production of double haploids can also be used for the development of inbreds in a breeding program. Double haploids are produced by the doubling of a set of chromosomes (1N) from a heterozygous plant to produce a completely homozygous individual. For example, see Wan *et al.*, "Efficient Production of Doubled Haploid Plants Through Colchicine Treatment of Anther-Derived Maize Callus", *Theoretical and Applied Genetics*, 77:889-892, 1989 and U.S. Application 2003/0005479. This can be advantageous because the process omits the generations of selfing needed to obtain a homozygous plant from a heterozygous source.

Hybrid seed production requires elimination or inactivation of pollen produced by the female parent. Incomplete removal or inactivation of the pollen provides the potential for self-pollination. This inadvertently self-pollinated seed may be unintentionally harvested and packaged with hybrid seed. Also, because the male parent is grown next to the female parent in the field there is the very low probability that the male selfed seed could be unintentionally harvested and packaged with the hybrid seed. Once the seed from the hybrid bag is planted, it is possible to identify and select these self-pollinated plants. These self-pollinated plants will be genetically equivalent to one of the inbred lines used to produce the hybrid. Though the possibility of inbreds being included in hybrid seed bags exists, the occurrence is very low because much care is taken to avoid such inclusions. It is worth noting that hybrid seed is sold to growers for the production of grain and forage and not for breeding or seed production.

By an individual skilled in plant breeding, these inbred plants unintentionally included in commercial hybrid seed can be identified and selected due to their decreased vigor when compared to the hybrid. Inbreds are identified by their less vigorous appearance for vegetative and/or reproductive characteristics, including shorter plant height, small ear size, ear and kernel shape, cob color, or other characteristics.

Identification of these self-pollinated lines can also be accomplished through molecular marker analyses. See, "The Identification of Female Selves in Hybrid Maize:

A Comparison Using Electrophoresis and Morphology", Smith, J.S.C. and Wych, R.D., Seed Science and Technology 14, pp. 1-8 (1995), the disclosure of which is expressly incorporated herein by reference. Through these technologies, the homozygosity of the self pollinated line can be verified by analyzing allelic composition at various loci along the genome. Those methods allow for rapid identification of the invention disclosed herein. See also, "Identification of Atypical Plants in Hybrid Maize Seed by Postcontrol and Electrophoresis" Sarca, V. *et al.*, Probleme de Genetica Teoritica si Aplicata Vol. 20 (1) pp. 29-42.

Another form of commercial hybrid production involves the use of a mixture of male sterile hybrid seed and male pollinator seed. When planted, the resulting male sterile hybrid plants are pollinated by the pollinator plants. This method is primarily used to produce grain with enhanced quality grain traits, such as high oil, because desired quality grain traits expressed in the pollinator will also be expressed in the grain produced on the male sterile hybrid plant. In this method the desired quality grain trait does not have to be incorporated by lengthy procedures such as recurrent backcross selection into an inbred parent line. One use of this method is described in U.S. Patent Nos. 5,704,160 and 5,706,603.

There are many important factors to be considered in the art of plant breeding, such as the ability to recognize important morphological and physiological characteristics, the ability to design evaluation techniques for genotypic and phenotypic traits of interest, and the ability to search out and exploit the genes for the desired traits in new or improved combinations.

The objective of commercial maize hybrid line development resulting from a maize plant breeding program is to develop new inbred lines to produce hybrids that combine to produce high grain yields and superior agronomic performance. One of the primary traits breeders seek is yield. However, many other major agronomic traits are of importance in hybrid combination and have an impact on yield or otherwise provide superior performance in hybrid combinations. Such traits include percent grain moisture at harvest, relative maturity, resistance to stalk breakage, resistance to root lodging, grain quality, and disease and insect resistance. In addition, the lines

per se must have acceptable performance for parental traits such as seed yields, kernel sizes, pollen production, all of which affect ability to provide parental lines in sufficient quantity and quality for hybridization. These traits have been shown to be under genetic control and many if not all of the traits are affected by multiple genes.

A breeder uses various methods to help determine which plants should be selected from the segregating populations and ultimately which inbred lines will be used to develop hybrids for commercialization. In addition to the knowledge of the germplasm and other skills the breeder uses, a part of the selection process is dependent on experimental design coupled with the use of statistical analysis. Experimental design and statistical analysis are used to help determine which plants, which family of plants, and finally which inbred lines and hybrid combinations are significantly better or different for one or more traits of interest. Experimental design methods are used to assess error so that differences between two inbred lines or two hybrid lines can be more accurately determined. Statistical analysis includes the calculation of mean values, determination of the statistical significance of the sources of variation, and the calculation of the appropriate variance components. Either a five or one percent significance level is customarily used to determine whether a difference that occurs for a given trait is real or due to the environment or experimental error. One of ordinary skill in the art of plant breeding would know how to evaluate the traits of two plant varieties to determine if there is no significant difference between the two traits expressed by those varieties. For example, see Fehr, Walt, Principles of Cultivar Development, pp. 261-286 (1987) which is incorporated herein by reference. Mean trait values may be used to determine whether trait differences are significant, and preferably the traits are measured on plants grown under the same environmental conditions.

Combining ability of a line, as well as the performance of the line per se, is a factor in the selection of improved maize inbreds. Combining ability refers to a line's contribution as a parent when crossed with other lines to form hybrids. The hybrids formed for the purpose of selecting superior lines are designated test crosses. One way of measuring combining ability is by using breeding values. Breeding values are

based in part on the overall mean of a number of test crosses. This mean is then adjusted to remove environmental effects and it is adjusted for known genetic relationships among the lines.

Once such a line is developed its value to society is substantial since it is important to advance the germplasm base as a whole in order to maintain or improve traits such as yield, disease resistance, pest resistance and plant performance in extreme weather conditions.

SUMMARY OF THE INVENTION

According to the invention, there is provided a hybrid maize plant, and its parts designated as 33N56, produced by crossing two Pioneer Hi-Bred International, Inc. proprietary inbred maize lines GE02757986 and GE587351. These lines, deposited with the American Type Culture Collection, (ATCC), Manassas, Virginia 20110, have Accession Number _____ for GE GE02757986 and Accession Number _____ for GE GE587351. This invention thus relates to the hybrid seed 33N56, the hybrid plant and its parts produced from the seed, and variants, mutants and trivial modifications of hybrid maize 33N56. This invention also relates to methods for producing a maize plant containing in its genetic material one or more transgenes and to the transgenic maize plants and their parts produced by those methods. This invention further relates to methods for producing maize lines derived from hybrid maize 33N56 and to the maize lines derived by the use of those methods. This hybrid maize plant is characterized by good yield and superior tolerance to Head Smut.

DEFINITIONS

Certain definitions used in the specification are provided below. In order to provide a clear and consistent understanding of the specification and claims, including the scope to be given such terms, the following definitions are provided. NOTE: ABS is in absolute terms and %MN is percent of the mean for the experiments in which the inbred or hybrid was grown. PCT designates that the trait is

calculated as a percentage. %NOT designates the percentage of plants that did not exhibit trait. For example, STKLDG %NOT is the percentage of plants in a plot that were not stalk lodged. These designators will follow the descriptors to denote how the values are to be interpreted. Below are the descriptors used in the data tables included herein.

ABTSTK = ARTIFICIAL BRITTLE STALK. A count of the number of "snapped" plants per plot following machine snapping. A snapped plant has its stalk completely snapped at a node between the base of the plant and the node above the ear. Expressed as percent of plants that did not snap.

ADF = PERCENT ACID DETERGENT FIBER. The percent of dry matter that is acid detergent fiber in chopped whole plant forage.

ALLELE. Any of one or more alternative forms of a genetic sequence. In a diploid cell or organism, the two alleles of a given sequence typically occupy corresponding loci on a pair of homologous chromosomes.

ANTROT = ANTHRACNOSE STALK ROT (*Colletotrichum graminicola*). A 1 to 9 visual rating indicating the resistance to Anthracnose Stalk Rot. A higher score indicates a higher resistance.

BACKCROSSING. Process in which a breeder crosses a hybrid progeny line back to one of the parental genotypes one or more times.

BARPLT = BARREN PLANTS. The percent of plants per plot that were not barren (lack ears).

BREEDING. The genetic manipulation of living organisms.

BREEDING CROSS. A cross to introduce new genetic material into a plant for the development of a new variety. For example, one could cross plant A with plant B, wherein plant B would be genetically different from plant A. After the breeding cross, the resulting F1 plants could then be selfed or sibbed for one, two, three or more times (F1, F2, F3, etc.) until a new inbred variety is developed. For clarification, such new inbred varieties would be within a pedigree distance of one breeding cross of plants A and B. The process described above would be referred to as one breeding cycle.

BRTSTK = BRITTLE STALKS. This is a measure of the stalk breakage near the time of pollination, and is an indication of whether a hybrid or inbred would snap or break near the time of flowering under severe winds. Data are presented as percentage of plants that did not snap.

CELL. Cell as used herein includes a plant cell, whether isolated, in tissue culture or incorporated in a plant or plant part.

CLDTST = COLD TEST. The percent of plants that germinate under cold test conditions.

CLN = CORN LETHAL NECROSIS. Synergistic interaction of maize chlorotic mottle virus (MCMV) in combination with either maize dwarf mosaic virus (MDMV-A or MDMV-B) or wheat streak mosaic virus (WSMV). A 1 to 9 visual rating indicating the resistance to Corn Lethal Necrosis. A higher score indicates a higher resistance.

COMRST = COMMON RUST (*Puccinia sorghi*). A 1 to 9 visual rating indicating the resistance to Common Rust. A higher score indicates a higher resistance.

CP = PERCENT OF CRUDE PROTEIN. The percent of dry matter that is crude protein in chopped whole plant forage.

CROSS POLLINATION. A plant is cross pollinated if the pollen comes from a flower on a different plant from a different family or line. Cross pollination excludes sib and self pollination.

CROSS. As used herein, the term "cross" or "crossing" can refer to a simple X by Y cross, or the process of backcrossing, depending on the context.

D/D = DRYDOWN. This represents the relative rate at which a hybrid will reach acceptable harvest moisture compared to other hybrids on a 1 to 9 rating scale. A high score indicates a hybrid that dries relatively fast while a low score indicates a hybrid that dries slowly.

DIPERS = DIPLODIA EAR MOLD SCORES (*Diplodia maydis* and *Diplodia macrospora*). A 1 to 9 visual rating indicating the resistance to Diplodia Ear Mold. A higher score indicates a higher resistance.

DIPLOID PLANT PART. Refers to a plant part or cell that has the same diploid genotype as 33N56.

DIPROT = DIPLODIA STALK ROT SCORE. Score of stalk rot severity due to *Diplodia* (*Diplodia maydis*). Expressed as a 1 to 9 score with 9 being highly resistant.

DM = PERCENT OF DRY MATTER. The percent of dry material in chopped whole plant silage.

DRPEAR = DROPPED EARS. A measure of the number of dropped ears per plot and represents the percentage of plants that did not drop ears prior to harvest.

D/T = DROUGHT TOLERANCE. This represents a 1 to 9 rating for drought tolerance, and is based on data obtained under stress conditions. A high score indicates good drought tolerance and a low score indicates poor drought tolerance.

EARHT = EAR HEIGHT. The ear height is a measure from the ground to the highest placed developed ear node attachment and is measured in centimeters.

EARMOLD = GENERAL EAR MOLD. Visual rating (1 to 9 score) where a "1" is very susceptible and a "9" is very resistant. This is based on overall rating for ear mold of mature ears without determining the specific mold organism, and may not be predictive for a specific ear mold.

EARSZ = EAR SIZE. A 1 to 9 visual rating of ear size. The higher the rating the larger the ear size.

EBTSTK = EARLY BRITTLE STALK. A count of the number of "snapped" plants per plot following severe winds when the corn plant is experiencing very rapid vegetative growth in the V5-V8 stage. Expressed as percent of plants that did not snap.

ECB1LF = EUROPEAN CORN BORER FIRST GENERATION LEAF FEEDING (*Ostrinia nubilalis*). A 1 to 9 visual rating indicating the resistance to preflowering leaf feeding by first generation European Corn Borer. A higher score indicates a higher resistance.

ECB2IT = EUROPEAN CORN BORER SECOND GENERATION INCHES OF TUNNELING (*Ostrinia nubilalis*). Average inches of tunneling per plant in the stalk.

ECB2SC = EUROPEAN CORN BORER SECOND GENERATION (*Ostrinia nubilalis*). A 1 to 9 visual rating indicating post flowering degree of stalk breakage and other evidence of feeding by European Corn Borer, Second Generation. A higher score indicates a higher resistance.

ECBDPE = EUROPEAN CORN BORER DROPPED EARS (*Ostrinia nubilalis*). Dropped ears due to European Corn Borer. Percentage of plants that did not drop ears under second generation corn borer infestation.

EGRWTH = EARLY GROWTH. This is a measure of the relative height and size of a corn seedling at the 2-4 leaf stage of growth. This is a visual rating (1 to 9), with 1 being weak or slow growth, 5 being average growth and 9 being strong growth. Taller plants, wider leaves, more green mass and darker color constitute a higher score.

ELITE INBRED. An inbred that contributed desirable qualities when used to produce commercial hybrids. An elite inbred may also be used in further breeding for the purpose of developing further improved varieties.

ERTLDG = EARLY ROOT LODGING. Early root lodging is the percentage of plants that do not root lodge prior to or around anthesis; plants that lean from the vertical axis at an approximately 30 degree angle or greater would be counted as root lodged.

ERTLPN = EARLY ROOT LODGING. An estimate of the percentage of plants that do not root lodge prior to or around anthesis; plants that lean from the vertical axis at an approximately 30 degree angle or greater would be considered as root lodged.

ERTLSC = EARLY ROOT LODGING SCORE. Score for severity of plants that lean from a vertical axis at an approximate 30 degree angle or greater which typically results from strong winds prior to or around flowering recorded within 2 weeks of a wind event. Expressed as a 1 to 9 score with 9 being no lodging.

ESTCNT = EARLY STAND COUNT. This is a measure of the stand establishment in the spring and represents the number of plants that emerge on per plot basis for the inbred or hybrid.

EYESPT = EYE SPOT (*Kabatiella zeae* or *Aureobasidium zeae*). A 1 to 9 visual rating indicating the resistance to Eye Spot. A higher score indicates a higher resistance.

FUSERS = FUSARIUM EAR ROT SCORE (*Fusarium moniliforme* or *Fusarium subglutinans*). A 1 to 9 visual rating indicating the resistance to Fusarium ear rot. A higher score indicates a higher resistance.

GDU = Growing Degree Units. Using the Barger Heat Unit Theory, which assumes that maize growth occurs in the temperature range 50°F - 86°F and that temperatures outside this range slow down growth; the maximum daily heat unit accumulation is 36 and the minimum daily heat unit accumulation is 0. The seasonal accumulation of GDU is a major factor in determining maturity zones.

GDUSHD = GDU TO SHED. The number of growing degree units (GDUs) or heat units required for an inbred line or hybrid to have approximately 50 percent of the plants shedding pollen and is measured from the time of planting. Growing degree units are calculated by the Barger Method, where the heat units for a 24-hour period are:

$$\text{GDU} = \frac{(\text{Max. temp.} + \text{Min. temp.}) - 50}{2}$$

The highest maximum temperature used is 86°F and the lowest minimum temperature used is 50°F. For each inbred or hybrid it takes a certain number of GDUs to reach various stages of plant development.

GDUSLK = GDU TO SILK. The number of growing degree units required for an inbred line or hybrid to have approximately 50 percent of the plants with silk emergence from time of planting. Growing degree units are calculated by the Barger Method as given in GDU SHD definition.

GENOTYPE. Refers to the genetic constitution of a cell or organism.

GIBERS = GIBBERELLA EAR ROT (PINK MOLD) (*Gibberella zeae*). A 1 to 9 visual rating indicating the resistance to Gibberella Ear Rot. A higher score indicates a higher resistance.

GIBROT = GIBBERELLA STALK ROT SCORE. Score of stalk rot severity due to *Gibberella* (*Gibberella zeae*). Expressed as a 1 to 9 score with 9 being highly resistant.

GLFSPT = GRAY LEAF SPOT (*Cercospora zeae-maydis*). A 1 to 9 visual rating indicating the resistance to Gray Leaf Spot. A higher score indicates a higher resistance.

GOSWLT = GOSS' WILT (*Corynebacterium nebraskense*). A 1 to 9 visual rating indicating the resistance to Goss' Wilt. A higher score indicates a higher resistance.

GRNAPP = GRAIN APPEARANCE. This is a 1 to 9 rating for the general appearance of the shelled grain as it is harvested based on such factors as the color of harvested grain, any mold on the grain, and any cracked grain. High scores indicate good grain quality.

H/POP = YIELD AT HIGH DENSITY. Yield ability at relatively high plant densities on a 1 to 9 relative rating system with a higher number indicating the hybrid responds well to high plant densities for yield relative to other hybrids. A 1, 5, and 9 would represent very poor, average, and very good yield response, respectively, to increased plant density.

HCBLT = HELMINTHOSPORIUM CARBONUM LEAF BLIGHT (*Helminthosporium carbonum*). A 1 to 9 visual rating indicating the resistance to Helminthosporium infection. A higher score indicates a higher resistance.

HD SMT = HEAD SMUT (*Sphacelotheca reiliana*). This score indicates the percentage of plants not infected.

HSKCVR = HUSK COVER. A 1 to 9 score based on performance relative to key checks, with a score of 1 indicating very short husks, tip of ear and kernels showing; 5 is intermediate coverage of the ear under most conditions, sometimes with thin husk; and a 9 has husks extending and closed beyond the tip of the ear. Scoring can best be done near physiological maturity stage or any time during dry down until harvested.

INC D/A = GROSS INCOME (DOLLARS PER ACRE). Relative income per acre assuming drying costs of two cents per point above 15.5 percent harvest moisture and current market price per bushel.

INCOME/ACRE. Income advantage of hybrid to be patented over other hybrid on per acre basis.

INC ADV = GROSS INCOME ADVANTAGE. GROSS INCOME advantage of variety #1 over variety #2.

KSZDCD = KERNEL SIZE DISCARD. The percent of discard seed; calculated as the sum of discarded tip kernels and extra large kernels.

LINKAGE. Refers to a phenomenon wherein alleles on the same chromosome tend to segregate together more often than expected by chance if their transmission was independent.

LINKAGE DISEQUILIBRIUM. Refers to a phenomenon wherein alleles tend to remain together in linkage groups when segregating from parents to offspring, with a greater frequency than expected from their individual frequencies.

L/POP = YIELD AT LOW DENSITY. Yield ability at relatively low plant densities on a 1 to 9 relative system with a higher number indicating the hybrid responds well to low plant densities for yield relative to other hybrids. A 1, 5, and 9 would represent very poor, average, and very good yield response, respectively, to low plant density.

LRTL DG = LATE ROOT LODGING. Late root lodging is the percentage of plants that do not root lodge after anthesis through harvest; plants that lean from the vertical axis at an approximately 30 degree angle or greater would be counted as root lodged.

LRTL PN = LATE ROOT LODGING. Late root lodging is an estimate of the percentage of plants that do not root lodge after anthesis through harvest; plants that lean from the vertical axis at an approximately 30 degree angle or greater would be considered as root lodged.

LRTL SC = LATE ROOT LODGING SCORE. Score for severity of plants that lean from a vertical axis at an approximate 30 degree angle or greater which typically

results from strong winds after flowering. Recorded prior to harvest when a root-lodging event has occurred. This lodging results in plants that are leaned or "lodged" over at the base of the plant and do not straighten or "goose-neck" back to a vertical position. Expressed as a 1 to 9 score with 9 being no lodging.

MDMCPX = MAIZE DWARF MOSAIC COMPLEX (MDMV = Maize Dwarf Mosaic Virus and MCDV = Maize Chlorotic Dwarf Virus). A 1 to 9 visual rating indicating the resistance to Maize Dwarf Mosaic Complex. A higher score indicates a higher resistance.

MST = HARVEST MOISTURE. The moisture is the actual percentage moisture of the grain at harvest.

MSTADV = MOISTURE ADVANTAGE. The moisture advantage of variety #1 over variety #2 as calculated by: $\text{MOISTURE of variety \#2} - \text{MOISTURE of variety \#1} = \text{MOISTURE ADVANTAGE of variety \#1}$.

NLFBLT = NORTHERN LEAF BLIGHT (*Helminthosporium turcicum* or *Exserohilum turcicum*). A 1 to 9 visual rating indicating the resistance to Northern Leaf Blight. A higher score indicates a higher resistance.

OILT = GRAIN OIL. Absolute value of oil content of the kernel as predicted by Near-Infrared Transmittance and expressed as a percent of dry matter.

PEDIGREE DISTANCE = Relationship among generations based on their ancestral links as evidenced in pedigrees. May be measured by the distance of the pedigree from a given starting point in the ancestry.

PERCENT IDENTITY. Percent identity as used herein refers to the comparison of the alleles of two plants or lines as scored by matching loci. Percent identity is determined by comparing a statistically significant number of the loci of two plants or lines and scoring a match when the same two alleles are present at the same loci for each plant. For example, a percent identity of 90% between hybrid 33N56 and another plant means that the two plants have the same two alleles at 90% of their loci.

PERCENT SIMILARITY. Percent similarity as used herein refers to the comparison of the alleles of two plants or lines as scored by matching alleles.

Percent similarity is determined by comparing a statistically significant number of the loci of two plants or lines and scoring one allele match when the same allele is present at the same loci for each plant and two allele matches when the same two alleles are present at the same loci for each plant. A percent similarity of 90% between hybrid 33N56 and another plant means that the two plants have 90% matching alleles.

PLANT. As used herein, the term "plant" includes reference to an immature or mature whole plant, including a plant that has been detasseled or from which seed or grain has been removed. Seed or embryo that will produce the plant is also considered to be the plant.

PLANT PARTS. As used herein, the term "plant parts" includes leaves, stems, roots, seed, grain, embryo, pollen, ovules, flowers, ears, cobs, husks, stalks, root tips, anthers, pericarp, silk, tissue, cells and the like.

PLTHT = PLANT HEIGHT. This is a measure of the height of the plant from the ground to the tip of the tassel in centimeters.

POLSC = POLLEN SCORE. A 1 to 9 visual rating indicating the amount of pollen shed. The higher the score the more pollen shed.

POLWT = POLLEN WEIGHT. This is calculated by dry weight of tassels collected as shedding commences minus dry weight from similar tassels harvested after shedding is complete.

POP K/A = PLANT POPULATIONS. Measured as 1000s per acre.

POP ADV = PLANT POPULATION ADVANTAGE. The plant population advantage of variety #1 over variety #2 as calculated by $\text{PLANT POPULATION of variety \#2} - \text{PLANT POPULATION of variety \#1} = \text{PLANT POPULATION ADVANTAGE of variety \#1}$.

PRM = PREDICTED RELATIVE MATURITY. This trait, predicted relative maturity, is based on the harvest moisture of the grain. The relative maturity rating is based on a known set of checks and utilizes standard linear regression analyses and is also referred to as the Comparative Relative Maturity Rating System that is similar to the Minnesota Relative Maturity Rating System.

PRMSHD = A relative measure of the growing degree units (GDU) required to reach 50% pollen shed. Relative values are predicted values from the linear regression of observed GDU's on relative maturity of commercial checks.

PROT = GRAIN PROTEIN. Absolute value of protein content of the kernel as predicted by Near-Infrared Transmittance and expressed as a percent of dry matter.

RTLDDG = ROOT LODGING. Root lodging is the percentage of plants that do not root lodge; plants that lean from the vertical axis at an approximately 30 degree angle or greater would be counted as root lodged.

RTLADV = ROOT LODGING ADVANTAGE. The root lodging advantage of variety #1 over variety #2.

SCTGRN = SCATTER GRAIN. A 1 to 9 visual rating indicating the amount of scatter grain (lack of pollination or kernel abortion) on the ear. The higher the score the less scatter grain.

SDGVGR = SEEDLING VIGOR. This is the visual rating (1 to 9) of the amount of vegetative growth after emergence at the seedling stage (approximately five leaves). A higher score indicates better vigor.

SEL IND = SELECTION INDEX. The selection index gives a single measure of the hybrid's worth based on information for up to five traits. A maize breeder may utilize his or her own set of traits for the selection index. One of the traits that is almost always included is yield. The selection index data presented in the tables represent the mean value averaged across testing stations.

SIL DMP = SILAGE DRY MATTER. The percent of dry material in chopped whole plant silage.

SELF POLLINATION. A plant is self-pollinated if pollen from one flower is transferred to the same or another flower of the same plant.

SIB POLLINATION. A plant is sib-pollinated when individuals within the same family or line are used for pollination.

SLFBLT = SOUTHERN LEAF BLIGHT (*Helminthosporium maydis* or *Bipolaris maydis*). A 1 to 9 visual rating indicating the resistance to Southern Leaf Blight. A higher score indicates a higher resistance.

SOURST = SOUTHERN RUST (*Puccinia polysora*). A 1 to 9 visual rating indicating the resistance to Southern Rust. A higher score indicates a higher resistance.

STAGRN = STAY GREEN. Stay green is the measure of plant health near the time of black layer formation (physiological maturity). A high score indicates better late-season plant health.

STARCH = PERCENT OF STARCH. The percent of dry matter that is starch in chopped whole plant forage.

STDADV = STALK STANDING ADVANTAGE. The advantage of variety #1 over variety #2 for the trait STK CNT.

STKCNT = NUMBER OF PLANTS. This is the final stand or number of plants per plot.

STKLDG = STALK LODGING REGULAR. This is the percentage of plants that did not stalk lodge (stalk breakage) at regular harvest (when grain moisture is between about 20 and 30%) as measured by either natural lodging or pushing the stalks and determining the percentage of plants that break below the ear.

STKLDL = LATE STALK LODGING. This is the percentage of plants that did not stalk lodge (stalk breakage) at or around late season harvest (when grain moisture is between about 15 and 18%) as measured by either natural lodging or pushing the stalks and determining the percentage of plants that break below the ear.

STKLDS = STALK LODGING SCORE. A plant is considered as stalk lodged if the stalk is broken or crimped between the ear and the ground. This can be caused by any or a combination of the following: strong winds late in the season, disease pressure within the stalks, ECB damage or genetically weak stalks. This trait should be taken just prior to or at harvest. Expressed on a 1 to 9 scale with 9 being no lodging.

STLLPN = LATE STALK LODGING. This is the percent of plants that did not stalk lodge (stalk breakage or crimping) at or around late season harvest (when grain moisture is below 20%) as measured by either natural lodging or pushing the stalks and determining the percentage of plants that break or crimp below the ear.

STLPCN = STALK LODGING REGULAR. This is an estimate of the percentage of plants that did not stalk lodge (stalk breakage at regular harvest (when grain moisture is between about 20 and 30%) as measured by either natural lodging or pushing the stalks and determining the percentage of plants that break below the ear.

STRT = GRAIN STARCH. Absolute value of starch content of the kernel as predicted by Near-Infrared Transmittance and expressed as a percent of dry matter.

STWWLT = Stewart's Wilt (*Erwinia stewartii*). A 1 to 9 visual rating indicating the resistance to Stewart's Wilt. A higher score indicates a higher resistance.

TASBLS = TASSEL BLAST. A 1 to 9 visual rating was used to measure the degree of blasting (necrosis due to heat stress) of the tassel at the time of flowering. A 1 would indicate a very high level of blasting at time of flowering, while a 9 would have no tassel blasting.

TASSZ = TASSEL SIZE. A 1 to 9 visual rating was used to indicate the relative size of the tassel. The higher the rating the larger the tassel.

TAS WT = TASSEL WEIGHT. This is the average weight of a tassel (grams) just prior to pollen shed.

TDM/HA = TOTAL DRY MATTER PER HECTARE. Yield of total dry plant material in metric tons per hectare.

TEXEAR = EAR TEXTURE. A 1 to 9 visual rating was used to indicate the relative hardness (smoothness of crown) of mature grain. A 1 would be very soft (extreme dent) while a 9 would be very hard (flinty or very smooth crown).

TILLER = TILLERS. A count of the number of tillers per plot that could possibly shed pollen was taken. Data are given as a percentage of tillers: number of tillers per plot divided by number of plants per plot.

TST WT = TEST WEIGHT (UNADJUSTED). The measure of the weight of the grain in pounds for a given volume (bushel).

TSWADV = TEST WEIGHT ADVANTAGE. The test weight advantage of variety #1 over variety #2.

WIN M% = PERCENT MOISTURE WINS.

WIN Y% = PERCENT YIELD WINS.

YIELD = YIELD OF SILAGE. Yield in tons per acre at 30% dry matter.

YIELD BU/A = YIELD (BUSHELS/ACRE). Yield of the grain at harvest in bushels per acre adjusted to 15% moisture.

YLDADV = YIELD ADVANTAGE. The yield advantage of variety #1 over variety #2 as calculated by: YIELD of variety #1 - YIELD variety #2 = yield advantage of variety #1.

YLD SC = YIELD SCORE. A 1 to 9 visual rating was used to give a relative rating for yield based on plot ear piles. The higher the rating the greater visual yield appearance.

Definitions for Area of Adaptability

When referring to area of adaptability, such term is used to describe the location with the environmental conditions that would be well suited for this maize line. Area of adaptability is based on a number of factors, for example: days to maturity, insect resistance, disease resistance, and drought resistance. Area of adaptability does not indicate that the maize line will grow in every location within the area of adaptability or that it will not grow outside the area.

Central Corn Belt: Iowa, Illinois, Indiana

Drylands: non-irrigated areas of North Dakota, South Dakota, Nebraska, Kansas, Colorado and Oklahoma

Eastern U.S.: Ohio, Pennsylvania, Delaware, Maryland, Virginia, and West Virginia

North central U.S.: Minnesota and Wisconsin

Northeast: Michigan, New York, Vermont, and Ontario and Quebec Canada

Northwest U.S.: North Dakota, South Dakota, Wyoming, Washington, Oregon, Montana, Utah, and Idaho

South central U.S.: Missouri, Tennessee, Kentucky, Arkansas

Southeast U.S.: North Carolina, South Carolina, Georgia, Florida, Alabama, Mississippi, and Louisiana

Southwest U.S.: Texas, Oklahoma, New Mexico, Arizona

Western U.S.: Nebraska, Kansas, Colorado, and California

Maritime Europe: Northern France, Germany, Belgium, Netherlands and Austria

DETAILED DESCRIPTION OF THE INVENTION

Inbred maize lines are typically developed for use in the production of hybrid maize lines. Maize hybrids need to be highly homogeneous, heterozygous and reproducible to be useful as commercial hybrids. There are many analytical methods available to determine the heterozygous nature and the identity of these lines.

The oldest and most traditional method of analysis is the observation of phenotypic traits. The data is usually collected in field experiments over the life of the maize plants to be examined. Phenotypic characteristics most often observed are for traits associated with plant morphology, ear and kernel morphology, insect and disease resistance, maturity, and yield.

In addition to phenotypic observations, the genotype of a plant can also be examined. A plant's genotype can be used to identify plants of the same variety or a related variety. For example, the genotype can be used to determine the pedigree of a plant. There are many laboratory-based techniques available for the analysis, comparison and characterization of plant genotype; among these are Isozyme Electrophoresis, Restriction Fragment Length Polymorphisms (RFLPs), Randomly Amplified Polymorphic DNAs (RAPDs), Arbitrarily Primed Polymerase Chain Reaction (AP-PCR), DNA Amplification Fingerprinting (DAF), Sequence Characterized Amplified Regions (SCARs), Amplified Fragment Length Polymorphisms (AFLPs), Simple Sequence Repeats (SSRs) which are also referred to as Microsatellites, and Single Nucleotide Polymorphisms (SNPs).

Isozyme Electrophoresis and RFLPs as discussed in Lee, M., "Inbred Lines of Maize and Their Molecular Markers," *The Maize Handbook*, (Springer-Verlag, New York, Inc. 1994, at 423-432) incorporated herein by reference, have been widely used to determine genetic composition. Isozyme Electrophoresis has a relatively low number of available markers and a low number of allelic variants. RFLPs allow more

discrimination because they have a higher degree of allelic variation in maize and a larger number of markers can be found. Both of these methods have been eclipsed by SSRs as discussed in Smith *et al.*, "An evaluation of the utility of SSR loci as molecular markers in maize (*Zea mays* L.): comparisons with data from RFLPs and pedigree", *Theoretical and Applied Genetics* (1997) vol. 95 at 163-173 and by Pejic *et al.*, "Comparative analysis of genetic similarity among maize inbreds detected by RFLPs, RAPDs, SSRs, and AFLPs," *Theoretical and Applied Genetics* (1998) at 1248-1255 incorporated herein by reference. SSR technology is more efficient and practical to use than RFLPs; more marker loci can be routinely used and more alleles per marker locus can be found using SSRs in comparison to RFLPs. Single Nucleotide Polymorphisms may also be used to identify the unique genetic composition of the invention and progeny lines retaining that unique genetic composition. Various molecular marker techniques may be used in combination to enhance overall resolution.

Maize DNA molecular marker linkage maps have been rapidly constructed and widely implemented in genetic studies. One such study is described in Boppenmaier, *et al.*, "Comparisons among strains of inbreds for RFLPs", *Maize Genetics Cooperative Newsletter*, 65:1991, pg. 90, is incorporated herein by reference.

Pioneer Brand Hybrid 33N56 is characterized by good yield, very good stalk strength, above average root lodging resistance and stay green. Hybrid 33N56 further demonstrates good test weight and good drought tolerance. The hybrid also demonstrates above average tolerance to Goss' Wilt, above average tolerance to Gray Leaf Spot, above average tolerance to Northern Leaf Blight, above average tolerance to Southern Leaf Blight, above average tolerance to Southern Rust, above average tolerance to Corn Lethal Necrosis, superior tolerance to Head Smut, above average tolerance to Diplodia Ear Rot and above average tolerance to Fusarium Ear and Kernel Rot. The hybrid is particularly suited to the Mediterranean and to the Eastern, South central, and Western areas of the United States.

Pioneer Brand Hybrid 33N56 is a single cross, yellow endosperm, dent flint maize hybrid. Hybrid 33N56 has a relative maturity of approximately 112 based on the Comparative Relative Maturity Rating System for harvest moisture of grain.

This hybrid has the following characteristics based on the data collected primarily at Johnston, Iowa.

TABLE 1
VARIETY DESCRIPTION INFORMATION
33N56

	AVG	STDEV	N
1. TYPE: (Describe intermediate types in comments section) 1=Sweet,2=Dent,3=Flint,4=Flour,5=Pop and 6=Ornamental. Comments: Dent-Flint	2		
2. MATURITY: DAYS HEAT UNITS	Days	H.Units	
Emergence to 50% of plants in silk	61	1,298	
Emergence to 50% of plants in pollen shed	61	1,308	
10% to 90% pollen shed	2	57	
50% Silk to harvest at 25% moisture			
3. PLANT:			
Plant Height (to tassel tip) (cm)	302.2	13.84	15
Ear Height (to base of top ear node) (cm)	117.7	8.13	15
Length of Top Ear Internode (cm)	18.7	2.58	15
Average Number of Tillers per Plant	0.1	0.05	3
Average Number of Ears per Stalk	1.0	0.02	3
Anthocyanin of Brace Roots: 1=Absent, 2=Faint, 3=Moderate, 4=Dark	3		
4. LEAF:			
Width of Ear Node Leaf (cm)	11.3	0.62	15
Length of Ear Node Leaf (cm)	92.7	5.37	15
Number of Leaves above Top Ear	7.6	0.51	15
Leaf Angle: (at anthesis, 2nd leaf above ear to stalk above leaf) (Degrees)	24.5	5.77	15
* Leaf Color: V. Dark Green Munsell: 5GY34			
Leaf Sheath Pubescence: 1=none to 9=like peach fuzz	4		
5. TASSEL:			
Number of Primary Lateral Branches	8.2	1.37	15
Branch Angle from Central Spike	22.1	10.59	15
Tassel Length: (from peduncle node to tassel tip),(cm).	62.8	4.13	15
Pollen Shed: 0=male sterile, 9=heavy shed	6		
* Anther Color: Pale Yellow Munsell: 10YR76			
* Glume Color: Light Green Munsell: 5GY76			
* Bar Glumes (glume bands): 1=absent, 2=present	1		
Peduncle Length: (from top leaf node to lower florets or branches),(cm).	20.1	4.07	15
* Munsell Glossy Book of Color, (A standard color reference). Kollmorgen Inst. Corp. New Windsor, NY.			

TABLE 1
VARIETY DESCRIPTION INFORMATION
33N56

Page 2

6a. EAR (Unhusked ear)	AVG	STDEV	N
* Silk color: Light Red Munsell: 5R58 (3 days after silk emergence)			
* Fresh husk color: Med. Green Munsell: 5GY78			
* Dry husk color: White Munsell: 10YR92 (65 days after 50% silking)			
Ear position at dry husk stage: 1=upright ,2=horizontal, 3=pendant	3		
Husk Tightness: (1=very loose, 9=very tight)	5		
Husk Extension (at harvest):1=short(ears exposed), 2=medium (<8cm), 3=long (8-10cm), 4=v. long (>10cm)	1		
6b. EAR (Husked ear data)			
Ear Length (cm):	17.7	0.88	15
Ear Diameter at mid-point (mm)	46.0	2.20	15
Ear Weight (gm):	172.2	23.05	15
Number of Kernel Rows:	16.7	2.09	15
Kernel Rows: 1=indistinct, 2=distinct	2		
Row Alignment: 1=straight, 2=slightly curved, 3=spiral	2		
Shank Length (cm):	8.5	1.85	15
Ear Taper: 1=slight cylind., 2=average, 3=extreme	2		
7. KERNEL (Dried):			
Kernel Length (mm):	11.9	0.92	15
Kernel Width (mm):	8.3	0.88	15
Kernel Thickness (mm):	4.1	0.35	15
Round Kernels (shape grade) (%)	12.6	5.07	3
Aleurone Color Pattern: 1=homozygous, 2=segregating	1		
* Aleurone Color: Yellow Munsell: 1.25Y816			
* Hard Endo. Color: Yellow Munsell: 10YR612			
Endosperm Type:	3		
1=sweet (su1), 2=extra sweet (sh2), 3=normal starch, 4=high amylose starch, 5=waxy starch, 6=high protein, 7=high lysine, 8=super sweet (se), 9=high oil, 10=other			
Weight per 100 Kernels (unsized sample) (gm):	25.3	1.53	3
8. COB:			
* Cob Diameter at mid-point (mm):	25.7	0.82	15
* Cob Color: Red Munsell: 2.5YR38			
* Munsell Glossy Book of color, (A standard color reference). Kollmorgen Inst. Corp. New Windsor, NY.			

TABLE 1
VARIETY DESCRIPTION INFORMATION
33N56

Page 3

10. DISEASE RESISTANCE:

(Rate from 1=most-susceptible to 9=most-resistant. Leave blank if not tested, leave race or strain options blank if polygenic.)

A. LEAF BLIGHTS, WILTS, AND LOCAL INFECTION DISEASES

Anthrachnose Leaf Blight (*Colletotrichum graminicola*)

Common Rust (*Puccinia sorghi*)

Common Smut (*Ustilago maydis*)

Eyespot (*Kabatiella zeae*)

8 Goss' Wilt (*Clavibacter michiganense* spp. *nebraskense*)

5 Gray Leaf Spot (*Cercospora zeae-maydis*)

Helminthosporium Leaf Spot (*Bipolaris zeicola*) Race:

6 Northern Leaf Blight (*Exserohilum turcicum*) Race:

5 Southern Leaf Blight (*Bipolaris maydis*) Race:

6 Southern Rust (*Puccinia polysora*)

Stewart's Wilt (*Erwinia stewartii*)

Other (Specify): _____

B. SYSTEMIC DISEASES

5 Corn Lethal Necrosis (MCMV and MDMV)

9 Head Smut (*Sphacelotheca reiliana*)

Maize Chlorotic Dwarf Virus (MDV)

Maize Chlorotic Mottle Virus (MCMV)

Maize Dwarf Mosaic Virus (MDMV)

Sorghum Downy Mildew of Corn (*Peronosclerospora sorghi*)

Other (Specify): _____

C. STALK ROTS

4 Anthracnose Stalk Rot (*Colletotrichum graminicola*)

Diplodia Stalk Rot (*Stenocarpella maydis*)

Fusarium Stalk Rot (*Fusarium moniliforme*)

Gibberella Stalk Rot (*Gibberella zeae*)

Other (Specify): _____

D. EAR AND KERNEL ROTS

Aspergillus Ear and Kernel Rot (*Aspergillus flavus*)

5 Diplodia Ear Rot (*Stenocarpella maydis*)

5 Fusarium Ear and Kernel Rot (*Fusarium moniliforme*)

Gibberella Ear Rot (*Gibberella zeae*)

Other (Specify): _____

TABLE 1
VARIETY DESCRIPTION INFORMATION
33N56

Page 4

11. INSECT RESISTANCE:

(Rate from 1=most-susceptible to 9=most-resist., leave blank if not tested.)

Corn Worm (*Helicoverpa zea*)

___ Leaf Feeding

___ Silk Feeding

___ Ear Damage

Corn Leaf Aphid (*Rhopalosiphum maydis*)

Corn Sap Beetle (*Capophilus dimidiatus*)

European Corn Borer (*Ostrinia nubilalis*)

1st. Generation (Typically whorl leaf feeding)

2nd. Generation (Typically leaf sheath-collar feeding)

___ Stalk Tunneling

___ cm tunneled/plant

Fall armyworm (*Spodoptera frugiperda*)

___ Leaf Feeding

___ Silk Feeding

___ mg larval wt.

Maize Weevil (*Sitophilus zeamais*)

Northern Rootworm (*Diabrotica barberi*)

Southern Rootworm (*Diabrotica undecimpunctata*)

Southwestern Corn Borer (*Diatraea grandiosella*)

___ Leaf Feeding

___ Stalk Tunneling

___ cm tunneled/plant

Two-spotted Spider Mite (*Tetranychus utricae*)

Western Rootworm (*Diabrotica virgifera virgifera*)

Other (Specify): _____

12. AGRONOMIC TRAITS:

5 Staygreen (at 65 days after anthesis; rate from 1-worst to 9-excellent)

% Dropped Ears (at 65 days after anthesis)

% Pre-anthesis Brittle Snapping

% Pre-anthesis Root Lodging

% Post-anthesis Root Lodging (at 65 days after anthesis)

% Post-anthesis Stalk Lodging

11,481.0 Kg/ha (Yield at 12-13% grain moisture)

Research Comparisons for Pioneer Hybrid 33N56

Comparisons of characteristics for Pioneer Brand Hybrid 33N56 were made against Hybrid 33B50, 33G26, 33M54, 33P66, 33K39 and 32K22.

Table 2A compares Pioneer Brand Hybrid 33N56 and Hybrid 33B50, a hybrid with a similar area of adaptation and similar maturity. The table demonstrates significant differences between Hybrid 33N56 and Hybrid 33B50 which include yield, number of growing degree units to pollen shed and to silk emergence, stalk count and resistance to Gray Leaf Spot.

Table 2B compares Pioneer Brand Hybrid 33N56 and Hybrid 33G26, a hybrid with similar maturity. Significant differences between Hybrid 33N56 and Hybrid 33G26 include yield, number of growing degree units to pollen shed and to silk emergence, stalk count, plant height and ear height.

Table 2C compares Pioneer Brand Hybrid 33N56 and Hybrid 33M54, a closely related hybrid with a similar area of adaptation. The table demonstrates significant differences between Hybrid 33N56 and Hybrid 33M54 which include harvest moisture, early stand count, growing degree units to pollen shed and to silk emergence, stalk count, resistance to Northern Leaf Blight, resistance to Diplodia Ear Mold and early root lodging resistance.

Table 2D compares Pioneer Brand Hybrid 33N56 and Hybrid 33P66, a hybrid with a similar area of adaptation. Significant differences between Hybrid 33N56 and Hybrid 33P66 include yield, harvest moisture, early growth score, growing degree units to pollen shed and to silk emergence, stalk count, plant height, ear height, stay green and resistance to Northern Leaf Blight.

Table 2E compares Pioneer Brand Hybrid 33N56 and Hybrid 33K39, a closely related hybrid with a similar area of adaptation. The table demonstrates significant differences between Hybrid 33N56 and Hybrid 33K39 which include harvest moisture, plant height and resistance to Northern Leaf Blight.

Table 2F compares Pioneer Brand Hybrid 33N56 and Hybrid 32K22, a closely related hybrid with a similar area of adaptation. Significant differences between Hybrid 33N56 and Hybrid 32K22 include harvest moisture, stalk count, plant height, ear height, resistance to Northern Leaf Blight and resistance to Diplodia Ear Mold.

TABLE 2A
HYBRID COMPARISON

Variety #1: 33N56

Variety #2: 33B50

Stat	YIELD BU/A 56# ABS	YIELD BU/A 56# %MN	MST PCT %MN	EGRWTH SCORE %MN	ESTCNT COUNT %MN	GDUSHD GDU %MN
Mean1	181.1	100.6	97.7	95.6	102.5	99.6
Mean2	174.3	97.1	97.9	90.6	98.5	95.4
Locs	111	111	113	16	11	23
Reps	122	122	124	18	14	30
Diff	6.8	3.5	0.2	5.0	3.9	4.3
Prob	0.005	0.008	0.780	0.437	0.074	0.000

Stat	GDUSLK GDU %MN	STKCNT COUNT %MN	PLTHT CM %MN	EARHT CM %MN	STAGR SCORE %MN	STKLDG %NOT %MN
Mean1	99.5	101.1	96.8	99.0	102.5	99.9
Mean2	96.0	99.9	91.0	87.5	93.2	99.6
Locs	19	189	32	30	30	1
Reps	25	275	38	32	33	1
Diff	3.5	1.1	5.8	11.5	9.3	0.3
Prob	0.000	0.001	0.000	0.000	0.305	

Stat	ABTSTK %NOT %MN	TSTWT LB/BU ABS	GLFSPT SCORE ABS	NLFBLT SCORE ABS	SLFBLT SCORE ABS	GOSWLT SCORE ABS
Mean1	93.5	56.6	4.8	5.5	4.9	8.0
Mean2	120.0	56.4	3.5	4.9	4.7	8.0
Locs	3	76	9	7	5	1
Reps	15	83	13	13	8	2
Diff	-26.5	0.2	1.3	0.6	0.2	0.0
Prob	0.063	0.430	0.037	0.222	0.772	

TABLE 2A (Continued)
HYBRID COMPARISON

Variety #1: 33N56

Variety #2: 33B50

Stat	ANTROT SCORE ABS	CLN SCORE ABS	FUSERS SCORE ABS	DIPERS SCORE ABS	SOURST SCORE ABS	ECBDPE %NOT ABS
Mean1	3.9	5.0	5.3	4.8	6.0	95.9
Mean2	4.0	4.7	5.4	4.2	3.0	98.6
Locs	6	1	6	3	1	1
Reps	12	3	10	6	1	1
Diff	-0.1	0.3	-0.2	0.7	3.0	-2.7
Prob	0.907		0.813	0.057		

Stat	ECB1LF SCORE ABS	ECB2SC SCORE ABS	HSKCVR SCORE ABS	GIBROT SCORE ABS	DIPROT SCORE ABS	BRTSTK %NOT ABS
Mean1	3.7	4.6	4.3	4.0	3.5	87.4
Mean2	3.7	4.3	5.4	2.7	7.0	99.8
Locs	1	4	12	3	1	4
Reps	3	8	12	6	2	6
Diff	0.0	0.3	-1.2	1.3	-3.5	-12.3
Prob		0.252	0.001	0.270		0.226

Stat	HDSMT %NOT ABS	ERTLPN %NOT ABS	LRTLPN %NOT ABS
Mean1	98.3	86.1	89.8
Mean2	94.5	88.3	91.5
Locs	5	9	16
Reps	9	10	18
Diff	3.8	-2.2	-1.7
Prob	0.200	0.169	0.771

TABLE 2B
HYBRID COMPARISON
Variety #1: 33N56
Variety #2: 33G26

Stat	YIELD BU/A 56# ABS	YIELD BU/A 56# %MN	MST PCT %MN	EGRWTH SCORE %MN	ESTCNT COUNT %MN	GDUSHD GDU %MN
Mean1	182.7	101.2	98.4	93.2	102.5	99.5
Mean2	173.5	96.4	95.4	108.6	99.5	101.8
Locs	155	155	156	19	11	35
Reps	168	168	169	21	15	42
Diff	9.2	4.8	-3.1	-15.4	3.0	-2.4
Prob	0.000	0.000	0.000	0.001	0.133	0.000

Stat	GDUSLK GDU %MN	STKCNT COUNT %MN	PLTHT CM %MN	EARHT CM %MN	STAGRN SCORE %MN	ERTLSC SCORE ABS
Mean1	99.2	101.2	97.2	97.6	104.0	7.0
Mean2	102.2	100.5	101.5	103.6	100.0	8.0
Locs	24	253	46	44	38	1
Reps	30	342	52	46	42	1
Diff	-3.0	0.7	-4.3	-6.0	4.0	-1.0
Prob	0.000	0.016	0.000	0.000	0.613	

Stat	STKLDS SCORE ABS	STKLDG %NOT %MN	ABTSTK %NOT %MN	DRPEAR %NOT %MN	TSTWT LB/BU ABS	GLFSPT SCORE ABS
Mean1	7.0	103.3	93.5	100.8	56.4	4.8
Mean2	6.9	97.8	137.6	100.8	57.2	5.7
Locs	13	2	3	1	106	9
Reps	13	2	15	1	115	13
Diff	0.1	5.5	-44.1	0.0	-0.8	-0.9
Prob	0.829	0.481	0.035		0.000	0.065

TABLE 2B (Continued)
HYBRID COMPARISON

Variety #1: 33N56

Variety #2: 33G26

Stat	NLFBLT SCORE ABS	SLFBLT SCORE ABS	GOSWLT SCORE ABS	ANTROT SCORE ABS	CLN SCORE ABS	FUSERS SCORE ABS
Mean1	5.5	4.9	8.0	4.4	5.0	5.3
Mean2	5.4	3.7	8.0	4.7	5.7	4.3
Locs	7	5	1	8	1	6
Reps	12	8	2	14	3	10
Diff	0.1	1.2	0.0	-0.3	-0.7	1.0
Prob	0.736	0.153		0.502		0.463

Stat	DIPERS SCORE ABS	SOURST SCORE ABS	ECBDPE %NOT ABS	ECB1LF SCORE ABS	ECB2SC SCORE ABS	HSKCVR SCORE ABS
Mean1	4.8	6.0	95.9	3.7	5.7	4.4
Mean2	4.0	5.0	97.3	4.3	5.5	4.3
Locs	3	1	1	1	8	11
Reps	5	1	1	3	12	11
Diff	0.8	1.0	-1.4	-0.7	0.2	0.1
Prob	0.130				0.593	0.588

Stat	GIBROT SCORE ABS	DIPROT SCORE ABS	BRTSTK %NOT ABS	HDSMT %NOT ABS	ERTLPN %NOT ABS	LRTLPN %NOT ABS
Mean1	4.0	3.5	83.3	98.3	87.5	90.9
Mean2	4.2	5.5	99.7	95.3	82.0	80.9
Locs	3	1	3	5	10	20
Reps	6	2	5	9	11	23
Diff	-0.2	-2.0	-16.4	3.1	5.5	9.9
Prob	0.840		0.239	0.159	0.360	0.158

TABLE 2C
HYBRID COMPARISON

Variety #1: 33N56

Variety #2: 33M54

Stat	YIELD BU/A 56# ABS	YIELD BU/A 56# %MN	MST PCT %MN	EGRWTH SCORE %MN	ESTCNT COUNT %MN	GDUSHD GDU %MN
Mean1	178.9	100.2	97.8	95.6	102.5	99.6
Mean2	178.1	99.8	103.9	93.3	96.0	100.8
Locs	110	110	110	16	11	23
Reps	120	120	120	18	15	30
Diff	0.7	0.4	6.0	2.4	6.5	-1.2
Prob	0.765	0.774	0.000	0.729	0.012	0.007

Stat	GDUSLK GDU %MN	STKCNT COUNT %MN	PLTHT CM %MN	EARHT CM %MN	STAGR SCORE %MN	STKLDG %NOT %MN
Mean1	99.5	101.1	96.9	99.0	102.3	99.9
Mean2	100.8	99.3	97.3	94.2	106.7	104.0
Locs	19	189	32	30	29	1
Reps	25	277	38	32	32	1
Diff	-1.3	1.8	-0.4	4.8	-4.4	-4.1
Prob	0.024	0.000	0.606	0.018	0.533	

Stat	ABTSTK %NOT %MN	TSTWT LB/BU ABS	GLFSPT SCORE ABS	NLFBLT SCORE ABS	SLFBLT SCORE ABS	GOSWLT SCORE ABS
Mean1	93.5	56.4	4.8	5.5	4.9	8.0
Mean2	102.7	57.4	5.9	4.5	7.0	8.0
Locs	3	74	9	7	5	1
Reps	15	80	13	13	8	2
Diff	-9.2	-1.0	-1.1	1.0	-2.1	0.0
Prob	0.015	0.000	0.048	0.004	0.003	

TABLE 2C (Continued)
HYBRID COMPARISON

Variety #1: 33N56

Variety #2: 33M54

Stat	ANTROT SCORE ABS	CLN SCORE ABS	FUSERS SCORE ABS	DIPERS SCORE ABS	SOURST SCORE ABS	ECBDPE %NOT ABS
Mean1	3.9	5.0	5.3	4.8	6.0	95.9
Mean2	3.7	5.0	6.3	3.5	3.0	95.9
Locs	6	1	6	3	1	1
Reps	10	3	10	6	1	1
Diff	0.3	0.0	-1.1	1.3	3.0	0.0
Prob	0.597		0.071	0.015		

Stat	ECB1LF SCORE ABS	ECB2SC SCORE ABS	HSKCVR SCORE ABS	GIBROT SCORE ABS	DIPROT SCORE ABS	BRTSTK %NOT ABS
Mean1	3.7	4.6	4.4	4.0	3.5	87.4
Mean2	4.0	4.7	5.8	4.7	3.0	98.9
Locs	1	4	11	3	1	4
Reps	3	8	11	6	2	6
Diff	-0.3	-0.1	-1.5	-0.7	0.5	-11.4
Prob		0.931	0.000	0.270		0.280

Stat	HDSMT %NOT ABS	ERTLPN %NOT ABS	LRTLPN %NOT ABS
Mean1	98.3	88.6	90.9
Mean2	89.2	68.6	88.7
Locs	5	11	18
Reps	9	14	21
Diff	9.2	20.0	2.3
Prob	0.090	0.037	0.633

TABLE 2D
HYBRID COMPARISON
Variety #1: 33N56
Variety #2: 33P66

Stat	YIELD BU/A 56# ABS	YIELD BU/A 56# %MN	MST PCT %MN	EGRWTH SCORE %MN	ESTCNT COUNT %MN	GDUSHD GDU %MN
Mean1	182.4	101.3	98.4	93.2	102.5	99.5
Mean2	178.3	99.1	100.2	79.8	101.0	100.8
Locs	156	156	157	19	11	35
Reps	168	168	169	21	15	42
Diff	4.0	2.1	1.7	13.4	1.5	-1.3
Prob	0.047	0.071	0.025	0.018	0.258	0.005

Stat	GDUSLK GDU %MN	STKCNT COUNT %MN	PLTHT CM %MN	EARHT CM %MN	STAGR SCORE %MN	ERTLSC SCORE ABS
Mean1	99.2	101.1	97.2	97.6	104.0	7.0
Mean2	100.5	99.2	98.8	99.9	76.2	4.0
Locs	24	256	46	44	38	1
Reps	30	346	52	46	41	1
Diff	-1.2	2.0	-1.6	-2.3	27.8	3.0
Prob	0.031	0.000	0.046	0.042	0.000	

Stat	STKLDS SCORE ABS	STKLDG %NOT %MN	ABTSTK %NOT %MN	DRPEAR %NOT %MN	TSTWT LB/BU ABS	GLFSPT SCORE ABS
Mean1	7.0	103.3	93.5	100.8	56.4	4.8
Mean2	7.2	81.2	90.3	100.8	56.7	4.5
Locs	13	2	3	1	108	9
Reps	13	2	15	1	116	13
Diff	-0.2	22.1	3.2	0.0	-0.4	0.3
Prob	0.760	0.216	0.896		0.024	0.347

TABLE 2D (Continued)
HYBRID COMPARISON

Variety #1: 33N56

Variety #2: 33P66

Stat	NLFBLT SCORE ABS	SLFBLT SCORE ABS	GOSWLT SCORE ABS	ANTROT SCORE ABS	CLN SCORE ABS	FUSERS SCORE ABS
Mean1	5.5	5.1	8.0	4.4	5.0	5.3
Mean2	4.4	4.8	8.0	4.5	4.0	5.8
Locs	7	4	1	8	1	6
Reps	13	7	2	14	3	10
Diff	1.1	0.4	0.0	-0.1	1.0	-0.6
Prob	0.027	0.391		0.871		0.239

Stat	DIPERS SCORE ABS	SOURST SCORE ABS	ECBDPE %NOT ABS	ECB1LF SCORE ABS	ECB2SC SCORE ABS	HSKCVR SCORE ABS
Mean1	4.8	6.0	95.9	3.7	5.7	4.3
Mean2	4.2	3.0	95.9	3.7	5.0	3.7
Locs	3	1	1	1	8	11
Reps	6	1	1	3	12	11
Diff	0.7	3.0	0.0	0.0	0.6	0.5
Prob	0.270				0.054	0.052

Stat	GIBROT SCORE ABS	DIPROT SCORE ABS	BRTSTK %NOT ABS	HDSMT %NOT ABS	ERTLPN %NOT ABS	LRTLPN %NOT ABS
Mean1	4.0	3.5	87.4	98.3	89.6	91.3
Mean2	2.8	2.0	98.7	98.0	90.4	86.6
Locs	3	1	4	5	12	21
Reps	6	2	6	9	15	25
Diff	1.2	1.5	-11.3	0.3	-0.8	4.7
Prob	0.250		0.274	0.835	0.795	0.374

TABLE 2E
HYBRID COMPARISON
Variety #1: 33N56
Variety #2: 33K39

Stat	YIELD BU/A 56# ABS	YIELD BU/A 56# %MN	MST PCT %MN	EGRWTH SCORE %MN	ESTCNT COUNT %MN	GDUSHD GDU %MN
Mean1	180.3	100.4	97.7	95.6	102.5	99.6
Mean2	184.9	103.2	99.5	108.1	102.4	100.0
Locs	115	115	115	16	11	23
Reps	128	128	128	18	15	30
Diff	-4.6	-2.9	1.8	-12.5	0.1	-0.4
Prob	0.044	0.032	0.025	0.017	0.960	0.388

Stat	GDUSLK GDU %MN	STKCNT COUNT %MN	PLTHT CM %MN	EARHT CM %MN	STAGR SCORE %MN	STKLDG %NOT %MN
Mean1	99.5	101.0	96.8	98.9	102.7	99.9
Mean2	99.5	100.5	99.5	101.0	104.2	96.7
Locs	19	195	33	31	29	1
Reps	25	288	39	33	33	1
Diff	0.0	0.5	-2.6	-2.2	-1.5	3.2
Prob	0.969	0.089	0.006	0.177	0.843	

Stat	ABTSTK %NOT %MN	TSTWT LB/BU ABS	GLFSPT SCORE ABS	NLFBLT SCORE ABS	SLFBLT SCORE ABS	GOSWLT SCORE ABS
Mean1	93.5	56.6	4.8	5.5	4.9	8.0
Mean2	93.3	57.4	5.8	3.6	4.3	8.0
Locs	3	78	9	7	5	1
Reps	15	87	13	13	8	2
Diff	0.2	-0.8	-0.9	1.9	0.6	0.0
Prob	0.991	0.000	0.055	0.002	0.236	

TABLE 2E (Continued)
HYBRID COMPARISON

Variety #1: 33N56

Variety #2: 33K39

Stat	ANTROT SCORE ABS	CLN SCORE ABS	FUSERS SCORE ABS	DIPERS SCORE ABS	SOURST SCORE ABS	ECBDPE %NOT ABS
Mean1	3.9	5.0	5.3	4.8	6.0	95.9
Mean2	4.1	5.0	4.7	4.5	5.0	95.9
Locs	6	1	6	3	1	1
Reps	12	3	10	6	1	1
Diff	-0.2	0.0	0.6	0.3	1.0	0.0
Prob	0.765		0.519	0.529		

Stat	ECB1LF SCORE ABS	ECB2SC SCORE ABS	HSKCVR SCORE ABS	GIBROT SCORE ABS	DIPROT SCORE ABS	BRTSTK %NOT ABS
Mean1	3.7	4.6	4.4	4.0	3.5	87.4
Mean2	3.3	4.1	4.9	3.8	5.5	98.0
Locs	1	4	11	3	1	4
Reps	3	8	11	6	2	6
Diff	0.3	0.5	-0.5	0.2	-2.0	-10.5
Prob		0.103	0.111	0.826		0.313

Stat	HDSMT %NOT ABS	ERTLPN %NOT ABS	LRTLPN %NOT ABS
Mean1	98.1	88.6	90.9
Mean2	88.7	85.9	85.9
Locs	7	11	18
Reps	13	14	22
Diff	9.4	2.7	5.0
Prob	0.055	0.441	0.325

TABLE 2F
HYBRID COMPARISON

Variety #1: 33N56

Variety #2: 32K22

Stat	YIELD BU/A 56# ABS	YIELD BU/A 56# %MN	MST PCT %MN	EGRWTH SCORE %MN	ESTCNT COUNT %MN	GDUSHD GDU %MN
Mean1	180.0	100.4	97.7	95.6	102.5	99.6
Mean2	188.8	105.7	106.2	101.9	103.3	99.2
Locs	116	116	116	16	11	23
Reps	129	129	129	18	15	30
Diff	-8.9	-5.4	8.5	-6.3	-0.8	0.4
Prob	0.000	0.000	0.000	0.236	0.758	0.402

Stat	GDUSLK GDU %MN	STKCNT COUNT %MN	PLTHT CM %MN	EARHT CM %MN	STAGR SCORE %MN	STKLDG %NOT %MN
Mean1	99.5	101.1	96.8	98.9	102.5	99.9
Mean2	99.8	100.3	100.5	103.1	118.1	100.8
Locs	19	194	33	31	30	1
Reps	25	284	39	33	34	1
Diff	-0.3	0.7	-3.7	-4.2	-15.5	-0.9
Prob	0.628	0.014	0.001	0.015	0.004	

Stat	ABTSTK %NOT %MN	TSTWT LB/BU ABS	GLFSPT SCORE ABS	NLFBLT SCORE ABS	SLFBLT SCORE ABS	GOSWLT SCORE ABS
Mean1	93.5	56.5	4.8	5.5	4.9	8.0
Mean2	109.6	56.1	4.8	4.1	5.2	7.0
Locs	3	80	9	7	5	1
Reps	14	89	13	12	8	2
Diff	-16.1	0.4	0.0	1.4	-0.3	1.0
Prob	0.134	0.072	1.000	0.023	0.305	

TABLE 2F (Continued)
HYBRID COMPARISON

Variety #1: 33N56

Variety #2: 32K22

Stat	ANTROT SCORE ABS	CLN SCORE ABS	FUSERS SCORE ABS	DIPERS SCORE ABS	SOURST SCORE ABS	ECBDPE %NOT ABS
Mean1	3.9	5.0	5.3	4.8	6.0	95.9
Mean2	4.8	4.0	5.5	4.0	5.0	98.6
Locs	6	1	6	3	1	1
Reps	12	3	10	6	1	1
Diff	-0.8	1.0	-0.3	0.8	1.0	-2.7
Prob	0.250		0.646	0.038		

Stat	ECB1LF SCORE ABS	ECB2SC SCORE ABS	HSKCVR SCORE ABS	GIBROT SCORE ABS	DIPROT SCORE ABS	BRTSTK %NOT ABS
Mean1	3.7	4.6	4.3	4.0	3.5	87.4
Mean2	4.3	4.8	6.1	3.5	2.5	98.8
Locs	1	4	12	3	1	4
Reps	3	8	12	6	2	6
Diff	-0.7	-0.2	-1.8	0.5	1.0	-11.4
Prob		0.391	0.000	0.478		0.245

Stat	HDSMT %NOT ABS	ERTLPN %NOT ABS	LRTLPN %NOT ABS
Mean1	98.3	88.6	90.9
Mean2	86.7	93.6	88.4
Locs	5	11	18
Reps	9	14	22
Diff	11.7	-5.0	2.6
Prob	0.115	0.587	0.358

Further Embodiments of the Invention

This invention also is directed to methods for producing a maize plant by crossing a first parent maize plant with a second parent maize plant wherein either the first or second parent maize plant is Pioneer Brand hybrid 33N56. In one embodiment the parent hybrid maize plant 33N56 will be crossed with another maize plant, sibbed, or selfed, to generate an inbred which may be used in the development of additional plants. In another embodiment, double haploid methods may be used to generate an inbred plant. Further, this invention is directed to methods for producing a hybrid 33N56-progeny maize plant by crossing hybrid maize plant 33N56 with itself or a second maize plant and growing the progeny seed, and repeating the crossing and growing steps with the hybrid maize 33N56-progeny plant from 1 to 2 times, 1 to 3 times, 1 to 4 times, or 1 to 5 times. Thus, any such methods using the hybrid maize plant 33N56 are part of this invention: selfing, sibbing, backcrosses, hybrid production, crosses to populations, and the like.

All plants produced using hybrid maize plant 33N56 as a parent are within the scope of this invention, including plants derived from hybrid maize plant 33N56. Progeny of the breeding methods described herein may be characterized in any number of ways, such as by traits retained in the progeny, pedigree and/or molecular markers. Combinations of these methods of characterization may be used. This includes varieties essentially derived from variety 33N56 with the term "essentially derived variety" having the meaning ascribed to such term in 7 U.S.C. § 2104(a)(3) of the Plant Variety Protection Act, which definition is hereby incorporated by reference. This also includes progeny plant and parts thereof with at least one ancestor that is hybrid maize plant 33N56 and more specifically where the pedigree of this progeny includes 1, 2, 3, 4, and/or 5 or cross pollinations to a maize plant 33N56, or a plant that has 33N56 as a progenitor. Pedigree is a method used by breeders of ordinary skill in the art to describe the varieties. Varieties that are more closely related by pedigree are likely to share common genotypes and combinations of phenotypic characteristics. All breeders of ordinary skill in the art maintain pedigree records of their breeding programs. These pedigree records contain a detailed description of

the breeding process, including a listing of all parental lines used in the breeding process and information on how such line was used. Thus, a breeder of ordinary skill in the art would know if 33N56 were used in the development of a progeny line, and would also know how many breeding crosses to a line other than 33N56 were made in the development of any progeny line. A progeny line so developed may then be used in crosses with other, different, maize inbreds to produce first generation (F_1) maize hybrid seeds and plants with superior characteristics.

Specific methods and products produced using hybrid maize plant in plant breeding are encompassed within the scope of the invention listed above. One such embodiment is the method of crossing hybrid maize plant 33N56 with itself to form a homozygous inbred parent line. Hybrid 33N56 would be sib or self pollinated to form a population of progeny plants. The population of progeny plants produced by this method is also an embodiment of the invention. This first population of progeny plants will have received all of its alleles from hybrid maize plant 33N56. The inbreeding process results in homozygous loci being generated and is repeated until the plant is homozygous at substantially every loci and becomes an inbred line. Once this is accomplished the inbred line may be used in crosses with other inbred lines, including but not limited to inbred parent lines disclosed herein to generate a first generation of F_1 hybrid plants. One of ordinary skill in the art can utilize breeder notebooks, or molecular methods to identify a particular hybrid plant produced using an inbred line derived from maize hybrid plant 33N56, in addition to comparing traits. Any such individual inbred plant is also encompassed by this invention.

These embodiments also include use of these methods with transgenic or backcross conversions of maize hybrid plant 33N56. Another such embodiment is a method of developing a line genetically similar to hybrid maize plant 33N56 in breeding that involves the repeated backcrossing of an inbred parent of, or an inbred line derived from, hybrid maize plant 33N56 to another different maize plant any number of times. Using backcrossing methods, or even the tissue culture and transgenic methods described herein, the backcross conversion methods described herein, or other breeding methods known to one of ordinary skill in the art, one can

develop individual plants, plant cells, and populations of plants that retain at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 99.5% genetic similarity or identity to maize hybrid plant 33N56. The percentage of the genetics retained in the progeny may be measured by either pedigree analysis or through the use of genetic techniques such as molecular markers or electrophoresis. In pedigree analysis, on average 50% of the starting germplasm would be passed to the progeny line after one cross to a different line, 25% after another cross to a different line, and so on. Molecular markers could also be used to confirm and/or determine the pedigree of the progeny line. The inbred parent would then be crossed to a second inbred parent of or derived from hybrid maize plant 33N56 to create hybrid maize plant 33N56 with additional beneficial traits such as transgenes or backcross conversions.

One method for producing a line derived from hybrid maize plant is as follows. One of ordinary skill in the art would obtain hybrid maize plant 33N56 and cross it with another variety of maize, such as an elite inbred variety. The F1 seed derived from this cross would be grown to form a population. The nuclear genome of the F1 would be made-up of 50% of hybrid maize plant 33N56 and 50% of the other elite variety. The F1 seed would be grown and allowed to self, thereby forming F2 seed. On average the F2 seed nuclear genome would have derived 50% of its alleles from the parent hybrid plant 33N56 and 50% from the other maize variety, but various individual plants from the population would have a much greater percentage of their alleles derived from the parent maize hybrid plant (Wang J. and R. Bernardo, 2000, Crop Sci. 40:659-665 and Bernardo, R. and A.L. Kahler, 2001, Theor. Appl. Genet 102:986-992). Molecular markers of 33N56, or its parents identified from routine screening of the deposited samples herein could be used to select and retain those lines with high similarity to 33N56. The F2 seed would be grown and selection of plants would be made based on visual observation, markers and/or measurement of traits. The traits used for selection may be any 33N56 trait described in this specification, including the hybrid maize plant 33N56 traits of yield, stalk strength,

root lodging resistance, stay green, test weight, tolerance to Goss' Wilt, tolerance to Gray Leaf Spot, tolerance to Northern Leaf Blight, tolerance to Southern Leaf Blight, tolerance to Southern Rust, tolerance to Corn Lethal Necrosis, tolerance to Head Smut, tolerance to Diplodia Ear Rot, tolerance to Fusarium Ear and Kernel Rot, and particularly suited to the Mediterranean and to the Eastern, South central, and Western areas of the United States.

Such traits may also be the good general or specific combining ability of 33N56, including its ability to produce backcross conversions, or other hybrids. The 33N56 progeny plants that exhibit one or more of the desired 33N56 traits, such as those listed above, would be selected and each plant would be harvested separately. This F3 seed from each plant would be grown in individual rows and allowed to self. Then selected rows or plants from the rows would be harvested individually. The selections would again be based on visual observation, markers and/or measurements for desirable traits of the plants, such as one or more of the desirable 33N56 traits listed above.

The process of growing and selection would be repeated any number of times until a 33N56 progeny plant is obtained. The 33N56 progeny inbred plant would contain desirable traits in hybrid combination derived from hybrid plant 33N56. The resulting progeny line would benefit from the efforts of the inventor(s), and would not have existed but for the inventor(s) work in creating 33N56. Another embodiment of the invention is a 33N56 progeny plant that has received the desirable 33N56 traits listed above through the use of 33N56, which traits were not exhibited by other plants used in the breeding process.

The previous example can be modified in numerous ways, for instance selection may or may not occur at every selfing generation, the hybrid may immediately be selfed without crossing to another plant, selection may occur before or after the actual self-pollination process occurs, or individual selections may be made by harvesting individual ears, plants, rows or plots at any point during the breeding process described. In addition, double haploid breeding methods may be used at any step in the process. The population of plants produced at each and any

cycle of breeding is also an embodiment of the invention, and on average each such population would predictably consist of plants containing approximately 50% of its genes from inbred parents of maize hybrid 33N56 in the first breeding cycle, 25% of its genes from inbred parents of maize hybrid 33N56 in the second breeding cycle, 12.5% of its genes from inbred parents of maize hybrid 33N56 in the third breeding cycle, 6.25% in the fourth breeding cycle, 3.125% in the fifth breeding cycle, and so on. In each case the use of 33N56 provides a substantial benefit. The linkage groups of 33N56 would be retained in the progeny lines, and since current estimates of the maize genome size is about 50,000-80,000 genes (Xiaowu, Gai *et al.*, Nucleic Acids Research, 2000, Vol. 28, No. 1, 94-96), in addition to a large amount of non-coding DNA that impacts gene expression, it provides a significant advantage to use 33N56 as starting material to produce a line that retains desired genetics or traits of 33N56.

Another embodiment of this invention is the method of obtaining a substantially homozygous 33N56 progeny plant by obtaining a seed from the cross of 33N56 and another maize plant and applying double haploid methods to the F1 seed or F1 plant or to any successive filial generation. Such methods substantially decrease the number of generations required to produce an inbred with similar genetics or characteristics to 33N56.

A further embodiment of the invention is a backcross conversion of 33N56 obtained by crossing inbred parent plants of hybrid maize plant 33N56, which comprise the backcross conversion. For a dominant or additive trait at least one of the inbred parents would include backcross conversion in its genome. For a recessive trait, each parent would include the backcross conversion in its genome. In each case the resultant hybrid maize plant 33N56 obtained from the cross of the parents includes a backcross conversion or transgene.

A backcross conversion of 33N56 occurs when DNA sequences are introduced through traditional (non-transformation) breeding techniques, such as backcrossing (Hallauer *et al.*, 1988), with a parent of 33N56 utilized as the recurrent parent. Both naturally occurring and transgenic DNA sequences may be introduced

through backcrossing techniques. The term backcross conversion is also referred to in the art as a single locus conversion. A backcross conversion may produce a plant with a trait or locus conversion in at least one or more backcrosses, including at least 2 crosses, at least 3 crosses, at least 4 crosses, at least 5 crosses and the like. Molecular marker assisted breeding or selection may be utilized to reduce the number of backcrosses necessary to achieve the backcross conversion. For example, see Openshaw, S.J. *et al.*, Marker-assisted Selection in Backcross Breeding. In: Proceedings Symposium of the Analysis of Molecular Data, August 1994, Crop Science Society of America, Corvallis, OR, where it is demonstrated that a backcross conversion can be made in as few as two backcrosses.

The complexity of the backcross conversion method depends on the type of trait being transferred (single genes or closely linked genes as vs. unlinked genes), the level of expression of the trait, the type of inheritance (cytoplasmic or nuclear) and the types of parents included in the cross. It is understood by those of ordinary skill in the art that for single gene traits that are relatively easy to classify, the backcross method is effective and relatively easy to manage. (See Hallauer *et al.* in Corn and Corn Improvement, Sprague and Dudley, Third Ed. 1998). Desired traits that may be transferred through backcross conversion include, but are not limited to, waxy starch, sterility (nuclear and cytoplasmic), fertility restoration, grain color (white), nutritional enhancements, drought tolerance, nitrogen utilization, altered fatty acid profile, increased digestibility, low phytate, industrial enhancements, disease resistance (bacterial, fungal or viral), insect resistance, herbicide resistance and yield enhancements. In addition, an introgression site itself, such as an FRT site, Lox site or other site specific integration site, may be inserted by backcrossing and utilized for direct insertion of one or more genes of interest into a specific plant variety. The trait of interest is transferred from the donor parent to the recurrent parent, in this case, an inbred parent of the maize plant disclosed herein. In some embodiments of the invention, the number of loci that may be backcrossed into 33N56 is at least 1, 2, 3, 4, or 5 and/or no more than 6, 5, 4, 3, or 2. A single loci may contain several transgenes, such as a transgene for disease resistance that, in the same expression

vector, also contains a transgene for herbicide resistance. The gene for herbicide resistance may be used as a selectable marker and/or as a phenotypic trait. A single locus conversion of site specific integration system allows for the integration of multiple genes at the converted loci.

The backcross conversion may result from either the transfer of a dominant allele or a recessive allele. Selection of progeny containing the trait of interest is accomplished by direct selection for a trait associated with a dominant allele. Transgenes transferred via backcrossing typically function as a dominant single gene trait and are relatively easy to classify. Selection of progeny for a trait that is transferred via a recessive allele, such as the waxy starch characteristic, requires growing and selfing the first backcross generation to determine which plants carry the recessive alleles. Recessive traits may require additional progeny testing in successive backcross generations to determine the presence of the locus of interest. The last backcross generation is usually selfed to give pure breeding progeny for the gene(s) being transferred, although a backcross conversion with a stably introgressed trait may also be maintained by further backcrossing to the recurrent parent with selection for the converted trait.

Along with selection for the trait of interest, progeny are selected for the phenotype of the recurrent parent. While occasionally additional polynucleotide sequences or genes may be transferred along with the backcross conversion, the backcross conversion line "fits into the same hybrid combination as the recurrent parent inbred line and contributes the effect of the additional gene added through the backcross." Poehlman *et al.* (1995, pg. 334). A progeny comprising at least 95%, 96%, 97%, 98%, 99%, 99.5% and 99.9% genetic identity to hybrid 33N56 and comprising the backcross conversion trait or traits of interest, is considered to be a backcross conversion of hybrid 33N56. It has been proposed that in general there should be at least four backcrosses when it is important that the recovered lines be essentially identical to the recurrent parent except for the characteristic being transferred (Fehr 1987, Principles of Cultivar Development). However, as noted above, the number of backcrosses necessary can be reduced with the use of

molecular markers. Other factors, such as a genetically similar donor parent, may also reduce the number of backcrosses necessary.

Hybrid seed production requires elimination or inactivation of pollen produced by the female inbred parent. Incomplete removal or inactivation of the pollen provides the potential for self-pollination. A reliable method of controlling male fertility in plants offers the opportunity for improved seed production. It should be understood that the plant can, through routine manipulation by detasseling, cytoplasmic genes, nuclear genes, or other factors, be produced in a male-sterile form. The term manipulated to be male sterile refers to the use of any available techniques to produce a male sterile version of maize line 33N56. The male sterility may be either partial or complete male sterility.

Hybrid maize seed is often produced by a male sterility system incorporating manual or mechanical detasseling. Alternate strips of two maize inbreds are planted in a field, and the pollen-bearing tassels are removed from one of the inbreds (female). Provided that there is sufficient isolation from sources of foreign maize pollen, the ears of the detasseled inbred will be fertilized only from the other inbred (male), and the resulting seed is therefore hybrid and will form hybrid plants.

Such embodiments are also within the scope of the present claims. This invention includes hybrid maize seed of 33N56 and the hybrid maize plant produced therefrom. The foregoing was set forth by way of example and is not intended to limit the scope of the invention.

This invention is also directed to the use of hybrid maize plant 33N56 in tissue culture. As used herein, the term plant includes plant protoplasts, plant cell tissue cultures from which maize plants can be regenerated, plant calli, plant clumps, and plant cells that are intact in plants, or parts of plants, such as embryos, pollen, ovules, flowers, kernels, ears, cobs, leaves, seeds, husks, stalks, roots, root tips, anthers, silk and the like. As used herein the phrase "growing the seed" or "grown from the seed" includes embryo rescue, isolation of cells from seed for use in tissue culture, as well as traditional growing methods.

Duncan, Williams, Zehr, and Widholm, Planta, (1985) 165:322-332 reflects that 97% of the plants cultured which produced callus were capable of plant regeneration. Subsequent experiments with both inbreds and hybrids produced 91% regenerable callus which produced plants. In a further study in 1988, Songstad, Duncan & Widholm in Plant Cell Reports (1988), 7:262-265 reports several media additions which enhance regenerability of callus of two inbred lines. Other published reports also indicated that "nontraditional" tissues are capable of producing somatic embryogenesis and plant regeneration. K.P. Rao, *et al.*, Maize Genetics Cooperation Newsletter, 60:64-65 (1986), refers to somatic embryogenesis from glume callus cultures and B.V. Conger, *et al.*, Plant Cell Reports, 6:345-347 (1987) indicates somatic embryogenesis from the tissue cultures of maize leaf segments. Thus, it is clear from the literature that the state of the art is such that these methods of obtaining plants are, and were, "conventional" in the sense that they are routinely used and have a very high rate of success.

Tissue culture of maize, including tassel/anther culture, is described in U.S. Application 2002/0062506A1 and European Patent Application, Publication No. 160,390, each of which are incorporated herein by reference. Maize tissue culture procedures are also described in Green and Rhodes, "Plant Regeneration in Tissue Culture of Maize," Maize for Biological Research (Plant Molecular Biology Association, Charlottesville, Virginia 1982, at 367-372) and in Duncan, *et al.*, "The Production of Callus Capable of Plant Regeneration from Immature Embryos of Numerous Zea Mays Genotypes," 165 Planta 322-332 (1985). Thus, another aspect of this invention is to provide cells which upon growth and differentiation produce maize plants having the genotype and/or physiological and morphological characteristics of hybrid maize plant 33N56.

The utility of hybrid maize plant 33N56 also extends to crosses with other species. Commonly, suitable species will be of the family Gramineae, and especially of the genera *Zea*, *Tripsacum*, *Coix*, *Schlerachne*, *Polytoca*, *Chionachne*, and *Trilobachne*, of the tribe Maydeae. Potentially suitable for crosses with 33N56 may be the various varieties of grain sorghum, *Sorghum bicolor* (L.) Moench.

Transformation of Maize

The advent of new molecular biological techniques has allowed the isolation and characterization of genetic elements with specific functions, such as encoding specific protein products. Scientists in the field of plant biology developed a strong interest in engineering the genome of plants to contain and express foreign genetic elements, or additional, or modified versions of native or endogenous genetic elements in order to alter the traits of a plant in a specific manner. Any DNA sequences, whether from a different species or from the same species, that are inserted into the genome using transformation are referred to herein collectively as "transgenes". In some embodiments of the invention, a transformed variant of 33N56 may contain at least one transgene but could contain at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 and/or no more than 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, or 2. Over the last fifteen to twenty years several methods for producing transgenic plants have been developed, and the present invention also relates to transformed versions of the claimed hybrid 33N56 as well as combinations thereof.

Numerous methods for plant transformation have been developed, including biological and physical plant transformation protocols. See, for example, Miki *et al.*, "Procedures for Introducing Foreign DNA into Plants" in *Methods in Plant Molecular Biology and Biotechnology*, Glick, B.R. and Thompson, J.E. Eds. (CRC Press, Inc., Boca Raton, 1993) pp. 67-88 and Armstrong, "The First Decade of Maize Transformation: A Review and Future Perspective" (*Maydica* 44:101-109, 1999). In addition, expression vectors and *in vitro* culture methods for plant cell or tissue transformation and regeneration of plants are available. See, for example, Gruber *et al.*, "Vectors for Plant Transformation" in *Methods in Plant Molecular Biology and Biotechnology*, Glick, B.R. and Thompson, J.E. Eds. (CRC Press, Inc., Boca Raton, 1993) pp. 89-119. See U.S. Patent No. 6,118,055, which is herein incorporated by reference.

The most prevalent types of plant transformation involve the construction of an expression vector. Such a vector comprises a DNA sequence that contains a gene

under the control of or operatively linked to a regulatory element, for example a promoter. The vector may contain one or more genes and one or more regulatory elements.

A genetic trait which has been engineered into the genome of a particular maize plant using transformation techniques, could be moved into the genome of another line using traditional breeding techniques that are well known in the plant breeding arts. These lines can then be crossed to generate a hybrid maize plant such as hybrid maize plant 33N56 which comprises a transgene. For example, a backcrossing approach is commonly used to move a transgene from a transformed maize plant to an elite inbred line, and the resulting progeny would then comprise the transgene(s). Also, if an inbred line was used for the transformation then the transgenic plants could be crossed to a different inbred in order to produce a transgenic hybrid maize plant. As used herein, "crossing" can refer to a simple X by Y cross, or the process of backcrossing, depending on the context.

Various genetic elements can be introduced into the plant genome using transformation. These elements include but are not limited to genes; coding sequences; inducible, constitutive, and tissue specific promoters; enhancing sequences; and signal and targeting sequences. For example, see the traits, genes and transformation methods listed in U.S. Patent No. 6,284,953, which is herein incorporated by reference.

With transgenic plants according to the present invention, a foreign protein can be produced in commercial quantities. Thus, techniques for the selection and propagation of transformed plants, which are well understood in the art, yield a plurality of transgenic plants that are harvested in a conventional manner, and a foreign protein then can be extracted from a tissue of interest or from total biomass. Protein extraction from plant biomass can be accomplished by known methods which are discussed, for example, by Heney and Orr, *Anal. Biochem.* 114: 92-6 (1981). In one embodiment, the biomass of interest is seed.

A genetic map can be generated, primarily via conventional Restriction Fragment Length Polymorphisms (RFLP), Polymerase Chain Reaction (PCR)

analysis, Simple Sequence Repeats (SSR) and Single Nucleotide Polymorphisms (SNP) which identifies the approximate chromosomal location of the integrated DNA molecule. For exemplary methodologies in this regard, see Glick and Thompson, METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY 269-284 (CRC Press, Boca Raton, 1993).

Wang *et al.* discuss "Large Scale Identification, Mapping and Genotyping of Single-Nucleotide Polymorphisms in the Human Genome", Science, 280:1077-1082, 1998, and similar capabilities are becoming increasingly available for the corn genome. Map information concerning chromosomal location is useful for proprietary protection of a subject transgenic plant. If unauthorized propagation is undertaken and crosses made with other germplasm, the map of the integration region can be compared to similar maps for suspect plants to determine if the latter have a common parentage with the subject plant. Map comparisons would involve hybridizations, RFLP, PCR, SSR and sequencing, all of which are conventional techniques. SNPs may also be used alone or in combination with other techniques.

Likewise, by means of the present invention, plants can be genetically engineered to express various phenotypes of agronomic interest. Through the transformation of maize the expression of genes can be modulated to enhance disease resistance, insect resistance, herbicide resistance, agronomic traits, grain quality and other traits. Transformation can also be used to insert DNA sequences which control or help control male-sterility. DNA sequences native to maize as well as non-native DNA sequences can be transformed into maize and used to modulate levels of native or non-native proteins. Various promoters, targeting sequences, enhancing sequences, and other DNA sequences can be inserted into the maize genome for the purpose of modulating the expression of proteins. Reduction of the activity of specific genes (also known as gene silencing, or gene suppression) is desirable for several aspects of genetic engineering in plants.

Many techniques for gene silencing are well known to one of skill in the art, including but not limited to antisense technology (see, *e.g.*, Sheehy *et al.* (1988) PNAS USA 85:8805-8809; and U.S. Patent Nos. 5,107,065; 5,453, 566; and

5,759,829); co-suppression (e.g., Taylor (1997) *Plant Cell* 9:1245; Jorgensen (1990) *Trends Biotech.* 8(12):340-344; Flavell (1994) *PNAS USA* 91:3490-3496; Finnegan *et al.* (1994) *Bio/Technology* 12: 883-888; and Neuhuber *et al.* (1994) *Mol. Gen. Genet.* 244:230-241); RNA interference (Napoli *et al.* (1990) *Plant Cell* 2:279-289; U.S. Patent No. 5,034,323; Sharp (1999) *Genes Dev.* 13:139-141; Zamore *et al.* (2000) *Cell* 101:25-33; and Montgomery *et al.* (1998) *PNAS USA* 95:15502-15507), virus-induced gene silencing (Burton, *et al.* (2000) *Plant Cell* 12:691-705; and Baulcombe (1999) *Curr. Op. Plant Bio.* 2:109-113); target-RNA-specific ribozymes (Haseloff *et al.* (1988) *Nature* 334: 585-591); hairpin structures (Smith *et al.* (2000) *Nature* 407:319-320; WO 99/53050; and WO 98/53083); ribozymes (Steinecke *et al.* (1992) *EMBO J.* 11:1525; and Perriman *et al.* (1993) *Antisense Res. Dev.* 3:253); oligonucleotide mediated targeted modification (e.g., WO 03/076574 and WO 99/25853); Zn-finger targeted molecules (e.g., WO 01/52620; WO 03/048345; and WO 00/42219); and other methods or combinations of the above methods known to those of skill in the art.

Exemplary transgenes useful for genetic engineering include, but are not limited to, those categorized below.

1. Transgenes That Confer Resistance To Pests or Disease And That Encode:
 - (A) Plant disease resistance genes. Plant defenses are often activated by specific interaction between the product of a disease resistance gene (R) in the plant and the product of a corresponding avirulence (Avr) gene in the pathogen. A plant variety can be transformed with cloned resistance gene to engineer plants that are resistant to specific pathogen strains. See, for example Jones *et al.*, *Science* 266: 789 (1994) (cloning of the tomato Cf-9 gene for resistance to *Cladosporium fulvum*); Martin *et al.*, *Science* 262: 1432 (1993) (tomato *Pto* gene for resistance to *Pseudomonas syringae* pv. tomato encodes a protein kinase); Mindrinos *et al.*, *Cell* 78: 1089 (1994) (*Arabidopsis RSP2* gene for resistance to *Pseudomonas syringae*). A plant resistant to a disease is one that is more resistant to a pathogen as compared to the wild type plant.

(B) A *Bacillus thuringiensis* protein, a derivative thereof or a synthetic polypeptide modeled thereon. See, for example, Geiser *et al.*, *Gene* 48: 109 (1986), who disclose the cloning and nucleotide sequence of a *Bt* delta-endotoxin gene. Moreover, DNA molecules encoding delta-endotoxin genes can be purchased from American Type Culture Collection (Rockville, MD), for example, under ATCC Accession Nos. 40098, 67136, 31995 and 31998. Other examples of *Bacillus thuringiensis* transgenes being genetically engineered are given in the following patents and hereby are incorporated by reference for this purpose: 5,188,960; 5,689,052; 5,880,275; WO 91/114778; WO 99/31248; WO 01/12731; WO 99/24581; WO 97/40162 and U.S. Patent Nos. 10/032,717; 10/414,637; and 10/606,320.

(C) An insect-specific hormone or pheromone such as an ecdysteroid and juvenile hormone, a variant thereof, a mimetic based thereon, or an antagonist or agonist thereof. See, for example, the disclosure by Hammock *et al.*, *Nature* 344: 458 (1990), of baculovirus expression of cloned juvenile hormone esterase, an inactivator of juvenile hormone.

(D) An insect-specific peptide or neuropeptide which, upon expression, disrupts the physiology of the affected pest. For example, see the disclosures of Regan, *J. Biol. Chem.* 269: 9 (1994) (expression cloning yields DNA coding for insect diuretic hormone receptor), and Pratt *et al.*, *Biochem. Biophys. Res. Comm.* 163: 1243 (1989) (an allostatin is identified in *Diploptera puntata*). See also U.S. Patent No. 5,266,317 to Tomalski *et al.*, who disclose genes encoding insect-specific, paralytic neurotoxins.

(E) An enzyme responsible for an hyperaccumulation of a monoterpenes, a sesquiterpene, a steroid, hydroxamic acid, a phenylpropanoid derivative or another non-protein molecule with insecticidal activity.

(F) An enzyme involved in the modification, including the post-translational modification, of a biologically active molecule; for example, a glycolytic enzyme, a proteolytic enzyme, a lipolytic enzyme, a nuclease, a cyclase, a transaminase, an esterase, a hydrolase, a phosphatase, a kinase, a phosphorylase, a polymerase, an elastase, a chitinase and a glucanase, whether natural or synthetic. See PCT

Application WO 93/02197 in the name of Scott *et al.*, which discloses the nucleotide sequence of a callase gene. DNA molecules which contain chitinase-encoding sequences can be obtained, for example, from the ATCC under Accession Nos. 39637 and 67152. See also Kramer *et al.*, *Insect Biochem. Molec. Biol.* 23: 691 (1993), who teach the nucleotide sequence of a cDNA encoding tobacco hookworm chitinase, and Kawalleck *et al.*, *Plant Molec. Biol.* 21: 673 (1993), who provide the nucleotide sequence of the parsley *ubi4-2* polyubiquitin gene.

(G) A molecule that stimulates signal transduction. For example, see the disclosure by Botella *et al.*, *Plant Molec. Biol.* 24: 757 (1994), of nucleotide sequences for mung bean calmodulin cDNA clones, and Griess *et al.*, *Plant Physiol.* 104: 1467 (1994), who provide the nucleotide sequence of a maize calmodulin cDNA clone.

(H) A hydrophobic moment peptide. See PCT Application WO 95/16776 (disclosure of peptide derivatives of Tachyplesin which inhibit fungal plant pathogens) and PCT Application WO 95/18855 (teaches synthetic antimicrobial peptides that confer disease resistance), the respective contents of which are hereby incorporated by reference for this purpose.

(I) A membrane permease, a channel former or a channel blocker. For example, see the disclosure by Jaynes *et al.*, *Plant Sci.* 89: 43 (1993), of heterologous expression of a cecropin-beta lytic peptide analog to render transgenic tobacco plants resistant to *Pseudomonas solanacearum*.

(J) A viral-invasive protein or a complex toxin derived therefrom. For example, the accumulation of viral coat proteins in transformed plant cells imparts resistance to viral infection and/or disease development effected by the virus from which the coat protein gene is derived, as well as by related viruses. See Beachy *et al.*, *Ann. Rev. Phytopathol.* 28: 451 (1990). Coat protein-mediated resistance has been conferred upon transformed plants against alfalfa mosaic virus, cucumber mosaic virus, tobacco streak virus, potato virus X, potato virus Y, tobacco etch virus, tobacco rattle virus and tobacco mosaic virus. *Id.*

(K) An insect-specific antibody or an immunotoxin derived therefrom. Thus,

an antibody targeted to a critical metabolic function in the insect gut would inactivate an affected enzyme, killing the insect. Cf. Taylor *et al.*, Abstract #497, SEVENTH INT'L SYMPOSIUM ON MOLECULAR PLANT-MICROBE INTERACTIONS (Edinburgh, Scotland, 1994) (enzymatic inactivation in transgenic tobacco via production of single-chain antibody fragments).

(L) A virus-specific antibody. See, for example, Tavladoraki *et al.*, *Nature* 366: 469 (1993), who show that transgenic plants expressing recombinant antibody genes are protected from virus attack.

(M) A developmental-arrestive protein produced in nature by a pathogen or a parasite. Thus, fungal endo alpha-1,4-D-polygalacturonases facilitate fungal colonization and plant nutrient release by solubilizing plant cell wall homo-alpha-1,4-D-galacturonase. See Lamb *et al.*, *Bio/Technology* 10: 1436 (1992). The cloning and characterization of a gene which encodes a bean endopolygalacturonase-inhibiting protein is described by Toubart *et al.*, *Plant J.* 2: 367 (1992).

(N) A developmental-arrestive protein produced in nature by a plant. For example, Logemann *et al.*, *Bio/Technology* 10: 305 (1992), have shown that transgenic plants expressing the barley ribosome-inactivating gene have an increased resistance to fungal disease.

(O) Genes involved in the Systemic Acquired Resistance (SAR) Response and/or the pathogenesis related genes. Briggs, S., *Current Biology*, 5(2) (1995).

(P) Antifungal genes (Cornelissen and Melchers, *Pl. Physiol.* 101:709-712, (1993) and Parijs *et al.*, *Planta* 183:258-264, (1991) and Bushnell *et al.*, *Can. J. of Plant Path.* 20(2):137-149 (1998).

(Q) Detoxification genes, such as for fumonisin, beauvericin, moniliformin and zearalenone and their structurally related derivatives. For example, see U.S. Patent No. 5,792,931.

(R) Cystatin and cysteine proteinase inhibitors.

(S) Defensin genes. See WO 03/000863.

(T) Genes conferring resistance to nematodes. See WO 03/033651 and Urwin *et al.*, *Planta* 204:472-479 (1998).

2. Transgenes That Confer Resistance To A Herbicide, For Example:

(A) A herbicide that inhibits the growing point or meristem, such as an imidazolinone or a sulfonylurea. Exemplary genes in this category code for mutant ALS and AHAS enzyme as described, for example, by Lee *et al.*, *EMBO J.* 7: 1241 (1988), and Miki *et al.*, *Theor. Appl. Genet.* 80: 449 (1990), respectively. See also, U.S. Patent Nos. 5,605,011; 5,013,659; 5,141,870; 5,767,361; 5,731,180; 5,304,732; 4,761,373; 5,331,107; 5,928,937; and 5,378,824; and international publication WO 96/33270, which are incorporated herein by reference in their entireties for this purpose.

(B) Glyphosate (resistance imparted by mutant 5-enolpyruvyl-3-phosphokimate synthase (EPSP), and *aroA* genes, respectively) and other phosphono compounds such as glufosinate (phosphinothricin acetyl transferase (PAT) and *Streptomyces hygroscopicus* phosphinothricin acetyl transferase (*bar*) genes), and pyridinoxy or phenoxy proprionic acids and cyclohexones (ACCase inhibitor-encoding genes). See, for example, U.S. Patent No. 4,940,835 to Shah *et al.*, which discloses the nucleotide sequence of a form of EPSPS which can confer glyphosate resistance. U.S. Patent No. 5,627,061 to Barry *et al.* also describes genes encoding EPSPS enzymes. See also U.S. Patent Nos. 6,248,876 B1; 6,040,497; 5,804,425; 5,633,435; 5,145,783; 4,971,908; 5,312,910; 5,188,642; 4,940,835; 5,866,775; 6,225,114 B1; 6,130,366; 5,310,667; 4,535,060; 4,769,061; 5,633,448; 5,510,471; Re. 36,449; RE 37,287 E; and 5,491,288; and international publications WO 97/04103; WO 97/04114; WO 00/66746; WO 01/66704; WO 00/66747 and WO 00/66748, which are incorporated herein by reference in their entireties for this purpose. Glyphosate resistance is also imparted to plants that express a gene that encodes a glyphosate oxido-reductase enzyme as described more fully in U.S. Patent Nos. 5,776,760 and 5,463,175, which are incorporated herein by reference in their entireties for this purpose. In addition glyphosate resistance can be imparted to plants by the over expression of genes encoding glyphosate N-acetyltransferase. See, for example, U.S. Application Serial Nos. 60/244,385; 60/377,175 and

60/377,719.

A DNA molecule encoding a mutant *aroA* gene can be obtained under ATCC Accession No. 39256, and the nucleotide sequence of the mutant gene is disclosed in U.S. Patent No. 4,769,061 to Comai. European Patent Application No. 0 333 033 to Kumada *et al.* and U.S. Patent No. 4,975,374 to Goodman *et al.* disclose nucleotide sequences of glutamine synthetase genes which confer resistance to herbicides such as L-phosphinothricin. The nucleotide sequence of a phosphinothricin-acetyl-transferase gene is provided in European Patent No. 0 242 246 and 0 242 236 to Leemans *et al.* De Greef *et al.*, *Bio/Technology* 7: 61 (1989), describe the production of transgenic plants that express chimeric *bar* genes coding for phosphinothricin acetyl transferase activity. See also, U.S. Patent Nos. 5,969,213; 5,489,520; 5,550,318; 5,874,265; 5,919,675; 5,561,236; 5,648,477; 5,646,024; 6,177,616 B1; and 5,879,903, which are incorporated herein by reference in their entireties for this purpose. Exemplary genes conferring resistance to phenoxy proprionic acids and cyclohexones, such as sethoxydim and haloxyfop, are the *Acc1-S1*, *Acc1-S2* and *Acc1-S3* genes described by Marshall *et al.*, *Theor. Appl. Genet.* 83: 435 (1992).

(C) A herbicide that inhibits photosynthesis, such as a triazine (*psbA* and *gs+* genes) and a benzonitrile (nitrilase gene). Przibilla *et al.*, *Plant Cell* 3: 169 (1991), describe the transformation of *Chlamydomonas* with plasmids encoding mutant *psbA* genes. Nucleotide sequences for nitrilase genes are disclosed in U.S. Patent No. 4,810,648 to Stalker, and DNA molecules containing these genes are available under ATCC Accession Nos. 53435, 67441 and 67442. Cloning and expression of DNA coding for a glutathione S-transferase is described by Hayes *et al.*, *Biochem. J.* 285: 173 (1992).

(D) Acetohydroxy acid synthase, which has been found to make plants that express this enzyme resistant to multiple types of herbicides, has been introduced into a variety of plants (see, e.g., Hattori *et al.* (1995) *Mol Gen Genet* 246:419). Other genes that confer tolerance to herbicides include: a gene encoding a chimeric protein of rat cytochrome P450A1 and yeast NADPH-cytochrome P450

oxidoreductase (Shiota *et al.* (1994) *Plant Physiol.* 106(1):17-23), genes for glutathione reductase and superoxide dismutase (Aono *et al.* (1995) *Plant Cell Physiol.* 36:1687, and genes for various phosphotransferases (Datta *et al.* (1992) *Plant Mol. Biol.* 20:619).

(E) Protoporphyrinogen oxidase (protox) is necessary for the production of chlorophyll, which is necessary for all plant survival. The protox enzyme serves as the target for a variety of herbicidal compounds. These herbicides also inhibit growth of all the different species of plants present, causing their total destruction. The development of plants containing altered protox activity which are resistant to these herbicides are described in U.S. Patent Nos. 6,288,306 B1; 6,282,837 B1; and 5,767,373; and international publication WO 01/12825, which are incorporated herein by reference in their entireties.

3. Transgenes That Confer Or Contribute To A Grain Trait, Such As:

(A) Modified fatty acid metabolism, for example, by

(1) Transforming a plant with an antisense gene of stearyl-ACP desaturase to increase stearic acid content of the plant. See Knultzon *et al.*, *Proc. Natl. Acad. Sci. USA* 89: 2624 (1992),

(2) Elevating oleic acid via FAD-2 gene modification and/or decreasing linolenic acid via FAD-3 gene modification (see U.S. Patent Nos. 6,063,947; 6,323,392; and WO 93/11245),

(3) Altering conjugated linolenic or linoleic acid content, such as in WO 01/12800,

(4) Modifying LEC1, AGP, Dek1, Superal1, thioredoxin, and/or a gamma zein knock out or mutant such as cs27 or TUSC 27. For example, see WO 02/42424, WO 98/22604, WO 03/011015, U.S. Patent No. 6,423,886 and Rivera-Madrid, R. *et al.*, *Proc. Natl. Acad. Sci.* 92:5620-5624 (1995).

(B) Decreased phytate content, for example, by the

(1) Introduction of a phytase-encoding gene would enhance breakdown of phytate, adding more free phosphate to the transformed plant. For

example, see Van Hartingsveldt *et al.*, *Gene* 127: 87 (1993), for a disclosure of the nucleotide sequence of an *Aspergillus niger* phytase gene.

(2) Introduction of a gene that reduces phytate content. In maize, this, for example, could be accomplished, by cloning and then re-introducing DNA associated with one or more of the alleles, such as the LPA alleles, identified in maize mutants characterized by low levels of phytic acid, such as in Raboy *et al.*, *Maydica* 35: 383 (1990) and/or by altering inositol kinase activity as in WO 02/059324, U.S. Application No. 2003/0009011, WO 03/027243, U.S. Application No. 2003/0079247 and WO 99/05298.

(C) Modified carbohydrate composition effected, for example, by transforming plants with a gene coding for an enzyme that alters the branching pattern of starch. See Shiroza *et al.*, *J. Bacteriol.* 170: 810 (1988) (nucleotide sequence of *Streptococcus mutans* fructosyltransferase gene), Steinmetz *et al.*, *Mol. Gen. Genet.* 200: 220 (1985) (nucleotide sequence of *Bacillus subtilis* levansucrase gene), Pen *et al.*, *Bio/Technology* 10: 292 (1992) (production of transgenic plants that express *Bacillus licheniformis* alpha-amylase), Elliot *et al.*, *Plant Molec. Biol.* 21: 515 (1993) (nucleotide sequences of tomato invertase genes), Sogaard *et al.*, *J. Biol. Chem.* 268: 22480 (1993) (site-directed mutagenesis of barley alpha-amylase gene), and Fisher *et al.*, *Plant Physiol.* 102: 1045 (1993) (maize endosperm starch branching enzyme II). The fatty acid modification genes mentioned above may also be used to effect starch content and/or composition through the interrelationship of the starch and oil pathways.

(D) Altered antioxidant content or composition, such as alteration of tocopherol or tocotrienols. For example, see WO 00/68393 involving the manipulation of antioxidant levels through alteration of a phytol prenyl transferase and WO 03/082899 through alteration of a homogentisate geranyl geranyl transferase.

(E) Improved digestibility and/or starch extraction through modification of UDP-D-xylose 4-epimerase, Fragile 1 and 2, Ref1, HCHL, C4H, such as in WO 99/10498.

4. Genes that Control Male-sterility:

(A) Introduction of a deacetylase gene under the control of a tapetum-specific promoter and with the application of the chemical N-Ac-PPT (WO 01/29237).

(B) Introduction of various stamen-specific promoters (WO 92/13956, WO 92/13957).

(C) Introduction of the barnase and the barstar gene (Paul *et al.*, *Plant Mol. Biol.* 19:611-622, 1992).

5. Genes that create a site for site specific DNA integration. This includes the introduction of FRT sites that may be used in the FLP/FRT system and/or Lox sites that may be used in the Cre/Loxp system. For example, see Lyznik, *et al.*, Site-Specific Recombination for Genetic Engineering in Plants, *Plant Cell Rep* (2003) 21:925-932 which is hereby incorporated by reference. Other systems that may be used include the Gin recombinase of phage Mu (Maeser *et al.*, 1991), the Pin recombinase of *E. coli* (Enomoto *et al.*, 1983), and the R/RS system of the pSR1 plasmid (Araki *et al.*, 1992).

6. Genes that affect growth characteristics, such as drought tolerance and nitrogen utilization. For example, see WO 00/73475 where water use efficiency is modulated through alteration of malate.

Genetic Marker Profile Through SSR

The present invention comprises a hybrid corn plant which is characterized by the molecular and physiological data presented herein and in the representative sample of said hybrid and of the inbred parents of said hybrid deposited with the ATCC.

To select and develop a superior hybrid, it is necessary to identify and select genetically unique individuals that occur in a segregating population. The segregating population is the result of a combination of crossover events plus the independent assortment of specific combinations of alleles at many gene loci that

results in specific and unique genotypes. Once such a line is developed its value to society is substantial since it is important to advance the germplasm base as a whole in order to maintain or improve traits such as yield, disease resistance, pest resistance and plant performance in extreme weather conditions. Backcross trait conversions are routinely used to add or modify one or a few traits of such a line and this further enhances its value and usefulness to society. The genetic variation among individual progeny of a breeding cross allows for the identification of rare and valuable new genotypes. Once identified, it is possible to utilize routine and predictable breeding methods to develop progeny that retain the rare and valuable new genotypes developed by the initial breeder.

Phenotypic traits exhibited by 33N56 can be used to characterize the genetic contribution of 33N56 to progeny lines developed through the use of 33N56. Quantitative traits including, but not limited to, yield, maturity, stay green, root lodging, stalk lodging, and early growth are typically governed by multiple genes at multiple loci. 33N56 progeny plants that retain the same degree of phenotypic expression of these quantitative traits as 33N56 have received significant genotypic and phenotypic contribution from 33N56. This characterization is enhanced when such quantitative trait is not exhibited in non-33N56 breeding material used to develop the 33N56 progeny.

As discussed, *supra*, in addition to phenotypic observations, a plant can also be described by its genotype. The genotype of a plant can be described through a genetic marker profile which can identify plants of the same variety, a related variety or be used to determine or validate a pedigree. Genetic marker profiles can be obtained by techniques such as Restriction Fragment Length Polymorphisms (RFLPs), Randomly Amplified Polymorphic DNAs (RAPDs), Arbitrarily Primed Polymerase Chain Reaction (AP-PCR), DNA Amplification Fingerprinting (DAF), Sequence Characterized Amplified Regions (SCARs), Amplified Fragment Length Polymorphisms (AFLPs), Simple Sequence Repeats (SSRs) which are also referred to as Microsatellites, and Single Nucleotide Polymorphisms (SNPs). For example, see Berry, Don, *et al.*, "Assessing Probability of Ancestry Using Simple Sequence

Repeat Profiles: Applications to Maize Hybrids and Inbreds", Genetics, 2002, 161:813-824, which is incorporated by reference herein in its entirety.

Particular markers used for these purposes are not limited to the set of markers disclosed herewithin, but are envisioned to include any type of marker and marker profile which provides a means of distinguishing varieties. In addition to being used for identification of inbred parents, hybrid variety 33N56, a hybrid produced through the use of 33N56 or its parents, and the identification or verification of pedigree for progeny plants produced through the use of 33N56, the genetic marker profile is also useful in breeding and developing backcross conversions.

Means of performing genetic marker profiles using SSR polymorphisms are well known in the art. SSRs are genetic markers based on polymorphisms in repeated nucleotide sequences, such as microsatellites. A marker system based on SSRs can be highly informative in linkage analysis relative to other marker systems in that multiple alleles may be present. Another advantage of this type of marker is that, through use of flanking primers, detection of SSRs can be achieved, for example, by the polymerase chain reaction (PCR), thereby eliminating the need for labor-intensive Southern hybridization. The PCRTM detection is done by use of two oligonucleotide primers flanking the polymorphic segment of repetitive DNA. Repeated cycles of heat denaturation of the DNA followed by annealing of the primers to their complementary sequences at low temperatures, and extension of the annealed primers with DNA polymerase, comprise the major part of the methodology.

Following amplification, markers can be scored by gel electrophoresis of the amplification products. Scoring of marker genotype is based on the size of the amplified fragment as measured by molecular weight (MW) rounded to the nearest integer. While variation in the primer used or in laboratory procedures can affect the reported molecular weight, relative values should remain constant regardless of the specific primer or laboratory used. When comparing lines it is preferable if all SSR profiles are performed in the same lab. An SSR service is available to the public on a contractual basis by DNA Landmarks in Saint-Jean-sur-Richelieu, Quebec, Canada (formerly Paragen, Celera AgGen, Perkin-Elmer AgGen, Linkage Genetics and NPI).

Primers used for the SSRs suggested herein are publicly available and may be found in the Maize GDB using the World Wide Web prefix followed by maizegdb.org (maintained by the USDA Agricultural Research Service), in Sharopova *et al.* (Plant Mol. Biol. 48(5-6):463-481), Lee *et al.* (Plant Mol. Biol. 48(5-6); 453-461). Primers may be constructed from publicly available sequence information. Some marker information may be available from DNA Landmarks.

A genetic marker profile of a hybrid should be the sum of its inbred parents, e.g., if one inbred parent is homozygous for allele x at a particular locus, and the other inbred parent is homozygous for allele y at that locus, the F1 hybrid will be x.y (heterozygous) at that locus. The profile can therefore be used to identify the inbred parents of hybrid 33N56. The determination of the male set of alleles and the female set of alleles may be made by profiling the hybrid and the pericarp of the hybrid seed, which is composed of maternal parent cells. The paternal parent profile is obtained by subtracting the pericarp profile from the hybrid profile.

In addition, plants and plant parts substantially benefiting from the use of 33N56 in their development such as 33N56 comprising a backcross conversion, transgene, or genetic sterility factor, may be identified by having a molecular marker profile with a high percent identity to 33N56. Such a percent identity might be 95%, 96%, 97%, 98%, 99%, 99.5% or 99.9% identical to 33N56.

The SSR profile of 33N56 also can be used to identify essentially derived varieties and other progeny lines developed from the use of 33N56, as well as cells and other plant parts thereof. Progeny plants and plant parts produced using 33N56 may be identified by having a molecular marker profile of at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 99.5% genetic contribution from hybrid maize plant 33N56.

Recurrent Selection and Mass Selection

Recurrent selection is a method used in a plant breeding program to improve a population of plants. 33N56 is suitable for use in a recurrent selection program. The

method entails individual plants cross pollinating with each other to form progeny. The progeny are grown and the superior progeny selected by any number of selection methods, which include individual plant, half-sib progeny, full-sib progeny, selfed progeny and topcrossing. The selected progeny are cross pollinated with each other to form progeny for another population. This population is planted and again superior plants are selected to cross pollinate with each other. Recurrent selection is a cyclical process and therefore can be repeated as many times as desired. The objective of recurrent selection is to improve the traits of a population. The improved population can then be used as a source of breeding material to obtain inbred lines to be used in hybrids or used as parents for a synthetic cultivar. A synthetic cultivar is the resultant progeny formed by the intercrossing of several selected inbreds.

Mass selection is a useful technique when used in conjunction with molecular marker enhanced selection. In mass selection seeds from individuals are selected based on phenotype and/or genotype. These selected seeds are then bulked and used to grow the next generation. Bulk selection requires growing a population of plants in a bulk plot, allowing the plants to self-pollinate, harvesting the seed in bulk and then using a sample of the seed harvested in bulk to plant the next generation. Instead of self pollination, directed pollination could be used as part of the breeding program.

Mutation Breeding

Mutation breeding is one of many methods that could be used to introduce new traits into 33N56. Mutations that occur spontaneously or are artificially induced can be useful sources of variability for a plant breeder. The goal of artificial mutagenesis is to increase the rate of mutation for a desired characteristic. Mutation rates can be increased by many different means including temperature, long-term seed storage, tissue culture conditions, radiation; such as X-rays, Gamma rays (e.g. cobalt 60 or cesium 137), neutrons, (product of nuclear fission by uranium 235 in an atomic reactor), Beta radiation (emitted from radioisotopes such as phosphorus 32 or carbon 14), or ultraviolet radiation (preferably from 2500 to 2900nm), or chemical

mutagens (such as base analogues (5-bromo-uracil), related compounds (8-ethoxy caffeine), antibiotics (streptonigrin), alkylating agents (sulfur mustards, nitrogen mustards, epoxides, ethylenamines, sulfates, sulfonates, sulfones, lactones), azide, hydroxylamine, nitrous acid, or acridines. Once a desired trait is observed through mutagenesis the trait may then be incorporated into existing germplasm by traditional breeding techniques, such as backcrossing. Details of mutation breeding can be found in "Principals of Cultivar Development" Fehr, 1993 Macmillan Publishing Company the disclosure of which is incorporated herein by reference. In addition, mutations created in other lines may be used to produce a backcross conversion of 33N56 that comprises such mutation.

Industrial Applicability

Maize is used as human food, livestock feed, and as raw material in industry. The food uses of maize, in addition to human consumption of maize kernels, include both products of dry- and wet-milling industries. The principal products of maize dry milling are grits, meal and flour. The maize wet-milling industry can provide maize starch, maize syrups, and dextrose for food use. Maize oil is recovered from maize germ, which is a by-product of both dry- and wet-milling industries.

Maize, including both grain and non-grain portions of the plant, is also used extensively as livestock feed, primarily for beef cattle, dairy cattle, hogs, and poultry.

Industrial uses of maize include production of ethanol, maize starch in the wet-milling industry and maize flour in the dry-milling industry. The industrial applications of maize starch and flour are based on functional properties, such as viscosity, film formation, adhesive properties, and ability to suspend particles. The maize starch and flour have application in the paper and textile industries. Other industrial uses include applications in adhesives, building materials, foundry binders, laundry starches, explosives, oil-well muds, and other mining applications.

Plant parts other than the grain of maize are also used in industry: for example, stalks and husks are made into paper and wallboard and cobs are used for fuel and to make charcoal.

The seed of the hybrid maize plant, the plant produced from the seed, a plant produced from crossing of maize hybrid plant 33N56 and various parts of the hybrid maize plant and transgenic versions of the foregoing, can be utilized for human food, livestock feed, and as a raw material in industry.

DEPOSITS

Applicant has made a deposit of at least 2500 seeds of hybrid maize 33N56 with the American Type Culture Collection (ATCC), Manassas, VA 20110 USA, ATCC Deposit No. _____. The seeds deposited with the ATCC on _____ were taken from the deposit maintained by Pioneer Hi-Bred International, Inc., 800 Capital Square, 400 Locust Street, Des Moines, Iowa 50309-2340 since prior to the filing date of this application. Access to this deposit will be available during the pendency of the application to the Commissioner of Patents and Trademarks and persons determined by the Commissioner to be entitled thereto upon request. Upon allowance of any claims in the application, the Applicant will make available to the public, pursuant to 37 C.F.R. § 1.808, sample(s) of the deposit of at least 2500 seeds of hybrid maize 33N56 with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, VA 20110-2209. This deposit of seed of hybrid maize 33N56 will be maintained in the ATCC depository, which is a public depository, for a period of 30 years, or 5 years after the most recent request, or for the enforceable life of the patent, whichever is longer, and will be replaced if it becomes nonviable during that period. Additionally, Applicant has satisfied all the requirements of 37 C.F.R. §§1.801 - 1.809, including providing an indication of the viability of the sample upon deposit. Applicant has no authority to waive any restrictions imposed by law on the transfer of biological material or its transportation in commerce. Applicant does not waive any infringement of their rights granted under this patent or under the Plant Variety Protection Act (7 USC 2321 et seq.). U.S. Plant Variety Protection of Hybrid Maize 33N56 has been applied for under Application No. _____.

All publications, patents and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All such publications, patents and patent applications are incorporated by reference herein to the same extent as if each was specifically and individually indicated to be incorporated by reference herein.

The foregoing invention has been described in detail by way of illustration and example for purposes of clarity and understanding. However, it will be obvious that certain changes and modifications such as backcross conversions and mutations, somoclonal variants, variant individuals selected from large populations of the plants of the instant inbred and the like may be practiced within the scope of the invention, as limited only by the scope of the appended claims.